

SECTION II

DRUG DOSAGE FORM AND DRUG DELIVERY SYSTEM DESIGN

CHAPTER

4

Dosage Form Design: Pharmaceutical and Formulation Considerations

OBJECTIVES

After reading this chapter, the student will be able to:

1. List reasons for the incorporation of drugs into various dosage forms
2. Compare and contrast the advantages/disadvantages of various drug dosage forms
3. Describe the information needed in preformulation studies to characterize a drug substance for possible inclusion into a dosage form
4. Describe the mechanisms of drug degradation and provide examples of each
5. Describe the five types of drug instability of concern to the practicing pharmacist
6. Summarize approaches employed to stabilize drugs in pharmaceutical dosage forms
7. Calculate rate reactions for various liquid dosage forms
8. Categorize various pharmaceutical ingredients and excipients

Drug substances are seldom administered alone; rather they are given as part of a formulation in combination with one or more nonmedicinal agents that serve varied and specialized pharmaceutical functions. Selective use of these nonmedicinal agents, referred to as pharmaceutical ingredients or excipients, produces dosage forms of various types. The pharmaceutical ingredients solubilize, suspend, thicken, dilute, emulsify, stabilize, preserve, color, flavor, and fashion medicinal agents into efficacious and appealing dosage forms. Each type of dosage form is unique in its physical and pharmaceutical characteristics. These varied preparations provide the manufacturing and compounding pharmacist with the challenges of formulation and the physician with the choice of drug and delivery system to

prescribe. The general area of study concerned with the formulation, manufacture, stability, and effectiveness of pharmaceutical dosage forms is termed pharmaceutics.

The proper design and formulation of a dosage form requires consideration of the physical, chemical, and biologic characteristics of all of the drug substances and pharmaceutical ingredients to be used in fabricating the product. The drug and pharmaceutical materials must be compatible with one another to produce a drug product that is stable, efficacious, attractive, easy to administer, and safe. The product should be manufactured with appropriate measures of quality control and packaged in containers that keep the product stable. The product should be labeled to promote correct use and be

stored under conditions that contribute to maximum shelf life.

Methods for the preparation of specific types of dosage forms and drug delivery systems are described in subsequent chapters. This chapter presents some general considerations regarding physical pharmacy, drug product formulation, and pharmaceutical ingredients.

THE NEED FOR DOSAGE FORMS

The potent nature and low dosage of most of the drugs in use today precludes any expectation that the general public could safely obtain the appropriate dose of a drug from the bulk material. Most drug substances are administered in milligram quantities, much too small to be weighed on anything but a sensitive prescription or electronic analytical balance. For instance, how could the lay person accurately obtain from a bulk supply the

325 mg of aspirin found in the common tablet? Not possible. Yet compared with many other drugs, the dose of aspirin is formidable (Table 4.1). For example, the dose of ethinyl estradiol, 0.05 mg, is 1/6,500 the amount of aspirin in an aspirin tablet. To put it another way, 6,500 ethinyl estradiol tablets, each containing 0.05 mg of drug, could be made from an amount of ethinyl estradiol equal to the amount of aspirin in just one standard tablet. When the dose of the drug is minute, as with ethinyl estradiol, solid dosage forms such as tablets and capsules must be prepared with fillers or diluents so that the dosage unit is large enough to pick up with the fingertips.

Besides providing the mechanism for the safe and convenient delivery of accurate dosage, dosage forms are needed for additional reasons:

- To protect the drug substance from the destructive influences of atmospheric oxygen or humidity (coated tablets, sealed ampuls)

TABLE 4.1 SOME DRUGS WITH RELATIVELY LOW USUAL DOSES

DRUG	USUAL DOSE (MG)	CATEGORY
Betaxolol HCl	10.00	Antianginal
Clotrimoxazole	10.00	Antifungal
Methylphenidate HCl	10.00	CNS stimulant
Medroxyprogesterone acetate	10.00	Progestin
Mesoridazine besylate	10.00	Antipsychotic
Morphine sulfate	10.00	Narcotic analgesic
Nifedipine	10.00	Coronary vasodilator
Omeprazole	10.00	Antiulcerative
Quinapril HCl	10.00	Antihypertensive
Chlorazepate dipotassium	7.50	Tranquilizer
Bupirone HCl	5.00	Antianxiety
Enalapril maleate	5.00	Antihypertensive
Hydrocodone	5.00	Narcotic analgesic
Prednisolone	5.00	Adrenocortical steroid
Albuterol sulfate	4.00	Bronchodilator
Chlorpheniramine maleate	4.00	Antihistaminic
Felodipine	2.50	Vasodilator
Glyburide	2.50	Antidiabetic
Doxazosin mesylate	2.00	Antihypertensive
Levorphanol tartrate	2.00	Narcotic analgesic
Prazosin HCl	2.00	Antihypertensive
Risperidone	2.00	Antipsychotic
Estropipate	1.25	Estrogen
Bumetanide	1.00	Diuretic
Clonazepam	1.00	Anticonvulsant
Ergoloid mesylates	1.00	Cognitive adjuvant
Alprazolam	0.50	Antianxiety
Colchicine	0.50	Gout suppressant
Nitroglycerin	0.40	Antianginal
Digoxin	0.25	Cardiotonic (maintenance)
Levothyroxine	0.10	Thyroid
Misoprostol	0.10	Antiulcerative, abortifacient
Ethinyl estradiol	0.05	Estrogen

- To protect the drug substance from the destructive influence of gastric acid after oral administration (enteric-coated tablets)
- To conceal the bitter, salty, or offensive taste or odor of a drug substance (capsules, coated tablets, flavored syrups)
- To provide liquid preparations of substances that are either insoluble or unstable in the desired vehicle (suspensions)
- To provide clear liquid dosage forms of substances (syrups, solutions)
- To provide rate-controlled drug action (various controlled-release tablets, capsules, and suspensions)
- To provide optimal drug action from topical administration sites (ointments, creams, transdermal patches, and ophthalmic, ear, and nasal preparations)
- To provide for insertion of a drug into one of the body's orifices (rectal or vaginal suppositories)
- To provide for placement of drugs directly in the bloodstream or body tissues (injections)
- To provide for optimal drug action through inhalation therapy (inhalants and inhalation aerosols)

GENERAL CONSIDERATIONS IN DOSAGE FORM DESIGN

Before formulating a drug substance into a dosage form, the desired product type must be determined insofar as possible to establish the framework for product development. Then, various initial formulations of the product are developed and examined for desired features (e.g., drug release profile, bioavailability, clinical effectiveness) and for pilot plant studies and production scale-up. The formulation that best meets the goals for the product is selected to be its *master formula*. Each batch of product subsequently prepared must meet the specifications established in the master formula.

There are many different forms into which a medicinal agent may be placed for the convenient and efficacious treatment of disease. Most commonly, a manufacturer prepares a drug substance in several dosage forms and strengths for the efficacious and convenient treatment of disease (Fig. 4.1). Before a medicinal agent is formulated into one or more dosage forms, among the factors considered are such therapeutic matters as the nature of the illness, the manner in which it is treated (locally or through systemic



FIGURE 4.1 Various forms of a drug substance marketed by a Pharmaceutical Company to meet the special requirements of the patient.

action), and the age and anticipated condition of the patient.

If the medication is intended for systemic use and oral administration is desired, tablets and/or capsules are usually prepared because they are easily handled by the patient and are most convenient in the self-administration of medication. If a drug substance has application in an emergency in which the patient may be comatose or unable to take oral medication, an injectable form of the medication may also be prepared. Many other examples of therapeutic situations affecting dosage form design could be cited, including motion sickness, nausea, and vomiting, for which tablets and skin patches are used for prevention and suppositories and injections for treatment.

The age of the intended patient also plays a role in dosage form design. For infants and children younger than 5 years of age, pharmaceutical liquids rather than solid forms are preferred for oral administration. These liquids, which are flavored aqueous solutions, syrups, or suspensions, are usually administered directly into the infant's or child's mouth by drop, spoon, or oral dispenser (Fig. 4.2) or incorporated into the child's food. A single liquid pediatric preparation may be used for infants and children of all ages, with the dose of the drug varied by the volume administered. When a young patient has a productive cough or is vomiting, gagging, or simply rebellious, there may be some question as to how much of the medicine administered is actually swallowed and how much is expectorated. In such instances, injections may be



FIGURE 4.2 Oral dosage devices to assist in measuring doses for children.

required. Infant-size rectal suppositories may also be employed, although drug absorption from the rectum is often erratic.

During childhood and even adulthood, a person may have difficulty swallowing solid dosage forms, especially uncoated tablets. For this reason, some medications are formulated as chewable tablets. Many of these tablets are comparable in texture to an after-dinner mint and break down into a pleasant-tasting creamy material. Newly available tablets dissolve in the mouth in about 10 to 15 seconds; this allows the patient to take a tablet but actually swallow a liquid. Capsules have been found by many to be more easily swallowed than whole tablets. If a capsule is moistened in the mouth before it is swallowed, it becomes slippery and readily slides down the throat with water. Also, a teaspoonful of gelatin dessert, liquid candy, or syrup placed in the mouth and partially swallowed before placing the solid dosage form in the mouth aids in swallowing them. Also, if a person has difficulty swallowing a capsule, the contents may be emptied into a spoon, mixed with jam, honey, or other similar food to mask the taste of the medication and swallowed. Medications intended for the elderly are commonly formulated into oral

liquids or may be extemporaneously prepared into an oral liquid by the pharmacist. However, certain tablets and capsules that are designed for controlled release should not be crushed or chewed, because that would interfere with their integrity and intended performance.

Many patients, particularly the elderly, take multiple medications daily. The more distinctive the size, shape, and color of solid dosage forms, the easier is proper identification of the medications. Errors in taking medications among the elderly occur frequently because of their multiple drug therapy and impaired eyesight. Dosage forms that allow reduced frequency of administration without sacrifice of efficiency are particularly advantageous.

In dealing with the problem of formulating a drug substance into a proper dosage form, research pharmacists employ knowledge gained through experience with other chemically similar drugs and through the proper use of the physical, chemical, biologic, and pharmaceutical sciences. The early stages of any new formulation include studies to collect basic information on the physical and chemical characteristics of the drug substance. These basic studies are the *preformulation* work needed before actual product formulation begins.

PREFORMULATION STUDIES

Before the formulation of a drug substance into a dosage form, it is essential that it be chemically and physically characterized. The following *preformulation studies* (1) and others provide the type of information needed to define the nature of the drug substance. This information provides the framework for the drug's combination with pharmaceutical ingredients in the fabrication of a dosage form.

Physical Description

It is important to understand the physical description of a drug substance prior to dosage form development. Most drug substances in use today are solid materials, pure chemical compounds of either crystalline or amorphous constitution. The purity of the chemical substance is essential for its identification and for evaluation of its chemical, physical, and biologic properties. Chemical properties include structure, form, and reactivity. Physical properties include such characteristics as its physical description, particle size, crystalline

structure, melting point, and solubility. Biologic properties relate to its ability to get to a site of action and elicit a biologic response.

Drugs can be used therapeutically as solids, liquids, and gases. Liquid drugs are used to a much lesser extent than solid drugs; gases, even less frequently.

Liquid drugs pose an interesting problem in the design of dosage forms and delivery systems. Many liquids are volatile and must be physically sealed from the atmosphere to prevent evaporation loss. Amyl nitrite, for example, is a clear yellowish liquid that is volatile even at low temperatures and is also highly flammable. It is kept for medicinal purposes in small sealed glass cylinders wrapped with gauze or another suitable material. When amyl nitrite is administered, the glass is broken between the fingertips, and the liquid wets the gauze covering, producing vapors that are inhaled by the patient requiring vasodilation. Propylhexedrine is another volatile liquid that must be contained in a closed system. This drug is used as a nasal inhalant for its vasoconstrictor action. A cylindrical roll of fibrous material is impregnated with propylhexedrine, and the saturated cylinder is placed in a suitable, usually plastic, sealed nasal inhaler. The inhaler's cap must be securely tightened each time it is used. Even then, the inhaler maintains its effectiveness for only a limited time because of the volatility of the drug.

Another problem associated with liquid drugs is that those intended for oral administration cannot generally be formulated into tablet form, the most popular form of oral medication, without chemical modification. An exception to this is the liquid drug nitroglycerin, which is formulated into sublingual tablets that disintegrate within seconds after placement under the tongue. However, because the drug is volatile, it has a tendency to escape from the tablets during storage, and it is critical that the tablets be stored in a tightly sealed glass container. For the most part, when a liquid drug is to be administered orally and a solid dosage form is desired, one of two approaches is used. First, the liquid substance may be sealed in a soft gelatin capsule. Vitamins A, D, and E, cyclosporin (Neoral, Sandimmune), and ergoloid mesylates (Hydergine LC) are liquids commercially available in capsule form. Second, the liquid drug may be developed into a solid ester or salt form that will be suitable for tablets or drug capsules. For instance, scopol-

amine hydrobromide is a solid salt of the liquid drug scopolamine and is easily pressed into tablets. Another approach to formulate liquids into solids is by mixing the drug with a solid or melted semisolid material, such as a high-molecular-weight polyethylene glycol. The melted mixture is poured into hard gelatin capsules to harden and the capsules sealed.

For certain liquid drugs, especially those taken orally in large doses or applied topically, their liquid nature may have some advantage in therapy. For example, 15-mL doses of mineral oil may be administered conveniently as such. Also, the liquid nature of undecylenic acid certainly does not hinder but rather enhances its use topically in the treatment of fungus infections of the skin. However, for the most part, pharmacists prefer solid materials in formulation work because they can easily form them into tablets and capsules.

Formulation and stability difficulties arise less frequently with solid dosage forms than with liquid preparations, and for this reason many new drugs first reach the market as tablets or dry-filled capsules. Later, when the pharmaceutical problems are resolved, a liquid form of the same drug may be marketed. This procedure is doubly advantageous, because for the most part physicians and patients alike prefer small, generally tasteless, accurately dosed tablets or capsules to the analogous liquid forms. Therefore, marketing a drug in solid form first is more practical for the manufacturer and suits most patients. It is estimated that tablets and capsules constitute the dosage form dispensed 70% of the time by community pharmacists, with tablets dispensed twice as frequently as capsules.

Microscopic Examination

Microscopic examination of the raw drug substance is an important step in preformulation work. It gives an indication of particle size and size range of the raw material along with the crystal structure. Photomicrographs of the initial and subsequent batch lots of the drug substance can provide important information in case of problems in formulation processing attributable to changes in particle or crystal characteristics of the drug. During some processing procedures, the solid drug powders must flow freely and not become entangled. Spherical and oval powders flow more easily than needle-shaped powders and make processing easier.

Heat of Vaporization

The use of vapor pressure is important in the operation of implantable pumps delivering medications as well as in aerosol dosage forms. Another application is the use of nasal inhalants (propylhexedrine with menthol and lavender oil-Benzedrex) for treating nasal congestion. In this latter dosage form, the quantity of drug required for effectiveness and a reasonable estimate of time of usefulness can be determined. Also, in the case of spills in inaccessible places, the time to evaporation of a substance can also be calculated. Some volatile drugs can even migrate within a tablet dosage form so the distribution may not be uniform any longer. This may have an impact in tablets that are scored for dosing where the drug in one portion may be higher or lower than in the other portion.

Exposure of personnel to hazardous drugs due to handling, spilling, or aerosolizing of drugs that may vaporize (oncology agents) is another application as the increase in mobility of the hazardous drug molecules may be related to temperature of the environment. Some drugs, such as carmustine, experience greater vapor pressures with increased temperature as compared

to cyclophosphamide, etoposide, cisplatin, and 5-fluorouracil, as illustrated in Physical Pharmacy Capsule 4.1, Heat of Vaporization.

Melting Point Depression

A characteristic of a pure substance is a defined melting point or melting range. If not pure, the substance will exhibit a change in melting point. This phenomenon is commonly used to determine the purity of a drug substance and in some cases the compatibility of various substances before inclusion in the same dosage form. This characteristic is further described in Physical Pharmacy Capsule 4.2, Melting Point Depression.

The Phase Rule

Phase diagrams are often constructed to provide a visual picture of the existence and extent of the presence of solid and liquid phases in binary, ternary, and other mixtures. Phase diagrams are normally two-component (binary) representations, as shown in Physical Pharmacy Capsule 4.3, The Phase Rule, but can also be three-component representations, as shown in Physical Pharmacy Capsule 4.4, Triangular Phase Diagram.



PHYSICAL PHARMACY CAPSULE 4.1

Heat of Vaporization

The amount of heat absorbed when 1 g of a liquid vaporizes is known as the heat of vaporization of that liquid and is measured in calories. The heat of vaporization of water at 100°C is 540 cal/g or about 9.720 cal/mole. This is the same quantity of heat energy that is released when 1 g of steam condenses to water at 100°C. This energy exchange is important in processes like steam sterilization as it is this energy transfer that results in death of microorganisms.

The movement of molecules varies with temperature. In liquids, this results in a tendency of the molecules to escape the liquid environment into a gaseous environment and possibly loss of the liquid. In the case of solids that sublime, the movement of the molecules is from the solid state to the vapor state. As an example, if one looks at an older bottle containing aspirin, there may be crystals of aspirin on the inside walls of the container. With ibuprofen, the walls of the container may become cloudy as the ibuprofen sublimates.

The use of vapor pressure is important in the operation of implantable pumps delivering medications as well as in aerosol dosage forms. Exposure of personnel to hazardous drugs due to handling, spilling, or aerosolizing of drugs that may vaporize (oncology agents) is another application as the increase in mobility of the hazardous drug molecules may be related to temperature of the environment. Some drugs, such as carmustine, experience greater vapor pressures with increased temperature as compared to cyclophosphamide, etoposide, cisplatin and 5-fluorouracil, as illustrated in the table below. Particle size affects vapor pressure; the smaller the particle size, the greater the vapor pressure. This demonstrates the importance of personnel protection with working with micronized hazardous powders. The time to evaporation of a substance can also be calculated.

PHYSICAL PHARMACY CAPSULE 4.1 CONT.

The variation of vapor pressure with temperature is described by the form of the Clausius–Clapeyron equation, as follows:

$$\frac{d \ln P}{dT} = \frac{\Delta H_{\text{vap}}}{RT^2}$$

assuming that ΔH_{vap} is constant, integration of the equation gives:

$$\log P = \frac{-\Delta H_{\text{vap}}}{2.303 RT} + \text{constant}$$

A plot of the log of the vapor pressure versus $1/T$ should be linear and the slope will equal $-\Delta H_{\text{vap}}/2.303R$ from which the enthalpy of vaporization can be calculated. With data obtained from Kiffmeyer TK, Kube C, Opiolka S, et al. Pharm J 2002;268:331, the following table can be constructed:

COMPOUND	MEASURED VAPOR PRESSURE (Pa)	
	20°C	40°C
Carmustine	0.019	0.530
Cisplatin	0.0018	0.0031
Cyclophosphamide	0.0033	0.0090
Etoposide	0.0026	0.0038
Fluorouracil	0.0014	0.0039

PHYSICAL PHARMACY CAPSULE 4.2

Melting Point Depression

The *melting point*, or *freezing point*, of a pure crystalline solid is defined as the temperature at which the pure liquid and solid exist in equilibrium. Drugs with a low melting point may soften during a processing step in which heat is generated, such as particle size reduction, compression, sintering, and so on. Also, the melting point or range of a drug can be used as an indicator of purity of chemical substances (a pure substance is ordinarily characterized by a very sharp melting peak). An altered peak or a peak at a different temperature may indicate an adulterated or impure drug. This is explained as follows.

The *latent heat of fusion* is the quantity of heat absorbed when 1 g of a solid melts; the molar heat of fusion (ΔH_f) is the quantity of heat absorbed when 1 mole of a solid melts. High-melting-point substances have high heat of fusion, and low-melting-point substances have low heat of fusion. These characteristics are related to the types of bonding in the specific substance. For example, ionic materials have high heats of fusion (NaCl melts at 801°C with a heat of fusion of 124 cal/g), and those with weaker van der Waals forces have low heats of fusion (paraffin melts at 52°C with a heat of fusion of 35.1 cal/g). Ice, with weaker hydrogen bonding, has a melting point of 0°C and a heat of fusion of 80 cal/g.

The addition of a second component to a pure compound (A), resulting in a mixture, will result in a melting point that is lower than that of the pure compound. The degree to which the melting point is lowered is proportional to the mole fraction (N_A) of the second component that is added. This can be expressed thus:

$$\Delta T = \frac{2.303 RTT_0}{\Delta H_f} \log N_A$$

where

ΔH_f is the molar heat of fusion,

T is the absolute equilibrium temperature,

T_0 is the melting point of pure A, and

R is the gas constant.

PHYSICAL PHARMACY CAPSULE 4.2 CONT.

Two noteworthy things contribute to the extent of lowering of the melting point:

1. Evident from this relationship is the inverse proportion between the melting point and the heat of fusion. When a second ingredient is added to a compound with a low molar heat of fusion, a large lowering of the melting point is observed; substances with a high molar heat of fusion will show little change in melting point with the addition of a second component.
2. The extent of lowering of the melting point is also related to the melting point itself. Compounds with low melting points are affected to a greater extent than compounds with high melting points upon the addition of a second component (i.e., low-melting-point compounds will result in a greater lowering of the melting point than those with high melting points).




PHYSICAL PHARMACY CAPSULE 4.3

The Phase Rule

A phase diagram, or temperature-composition diagram, represents the melting point as a function of composition of two or three component systems. The figure is an example of such a representation for a two-component mixture.

This phase diagram depicts a two-component mixture in which the components are completely miscible in the molten state and no solid solution or addition compound is formed in the solid state. As is evident, starting from the extremes of either pure component A or pure component B, as the second component is added, the melting point of the pure component decreases. There is a point on this phase diagram at which a minimum melting point occurs (i.e., the eutectic point). As is evident, four regions, or phases, in this diagram, represent the following:



- I Solid A + solid B
- II Solid A + melt
- III Solid B + melt
- IV Melt

Each phase is a homogenous part of the system, physically separated by distinct boundaries.

A description of the conditions under which these phases can exist is called the *Phase Rule*, which can be presented thus:

$$F = C - P + X$$

where

F is the number of degrees of freedom,

C is the number of components,

P is the number of phases, and

X is a variable dependent upon selected considerations of the phase diagram (1, 2, or 3).

C describes the minimum number of chemical components to be specified to define the phases. F is the number of independent variables that must be specified to define the complete system (e.g., temperature, pressure, concentration).

PHYSICAL PHARMACY CAPSULE 4.3 CONT.

EXAMPLE 1

In a mixture of menthol and thymol, a phase diagram similar to that illustrated can be obtained. To describe the number of degrees of freedom in the part of the graph moving from the curved line starting at pure A, progressing downward to the eutectic point, and then following an increasing melting point to pure B, it is evident from this presentation that either temperature or composition will describe this system, since it is assumed in this instance that pressure is constant. Therefore, the number of degrees of freedom to describe this portion of the phase diagram is given thus:

$$F = 2 - 2 + 1 = 1$$

In other words, along this line either temperature or composition will describe the system.

EXAMPLE 2

When in the area of a single phase of the diagram, such as the melt (IV), the system can be described thus:

$$F = 2 - 1 + 1 = 2$$

In this portion of the phase diagram, two factors, temperature and composition, can be varied without a change in the number of phases in the system.

EXAMPLE 3

At the eutectic point,

$$F = 2 - 3 + 1 = 0$$

and any change in the concentration or temperature may cause disappearance of one of the two solid phases or the liquid phase.

Phase diagrams are valuable for interpreting interactions between two or more components, relating not only to melting point depression and possible liquefaction at room temperature but also the formation of solid solutions, coprecipitates, and other solid-state interactions.

PHYSICAL PHARMACY CAPSULE 4.4

Triangular (Three-component) Phase Diagram


A three-component phase diagram has four degrees of freedom: $F = 3 - 1 + 2 = 4$. In this case, temperature and pressure are two of the conditions and the concentrations of two of the three components make up the rest. Only two concentrations are required because the third will be the difference between 100% and the sum of the other two components.

These systems are used for determining miscibility/solubility, coacervation regions, gel-forming regions for multicomponent mixtures, etc. To read a 3-phase diagram, each of the three corners of the triangle represent 100% by weight of one of the components (A, B, C) and 0% by weight of the other two (A, B, C). The lines joining the corner points forming the triangle each represent two component mixtures of the three possible combinations (AB, BC, and CA). If two of the components are known, the third is known by difference. Any combination of the three components is described by a single point on the diagram. Combining different proportions of the three components and observing for an end point (solubility, gel-formation, haziness, etc.), the phase differences can be visualized, as follows.

PHYSICAL PHARMACY CAPSULE 4.4 CONT.

The following is a stack of four separate pseudoternary phase diagrams for a quaternary system composed of Brij 96, glycerin, mineral oil, and water. The Brij 96:glycerin ratio is noted on the diagram and is considered one of three components. The shaded regions represent gel systems while the clear regions represent fluid systems.

In addition to observing the phase changes in a single plane, the use of stacked ternary phase diagrams enables one to visualize the change using different ratios of one of the components (in this case, the Brij 96:glycerin ratios). Constructions like this enable a pharmaceutical scientist to select the best ratios and combinations of components for a formulation.



Particle Size

Certain physical and chemical properties of drug substances, including dissolution rate, bioavailability, content uniformity, taste, texture, color, and stability, are affected by the particle size distribution. In addition, flow characteristics and sedimentation rates, among other properties, are important factors related to particle size. It is essential to establish as early as possible how the particle size of the drug substance may affect formulation and efficacy. Of special interest is the effect of particle size on absorption. Particle size significantly influences the oral absorption profiles of certain drugs, including griseofulvin, nitrofurantoin, spironolactone, and procaine penicillin. Also, satisfactory con-

tent uniformity in solid dosage forms depends to a large degree on particle size and the equal distribution of the active ingredient throughout the formulation. Particle size is discussed further in Chapter 6. Figure 4.3 shows a particle size analyzer.

Polymorphism

An important factor on formulation is the crystal or amorphous form of the drug substance. Polymorphic forms usually exhibit different physicochemical properties, including melting point and solubility. Polymorphic forms in drugs are relatively common. It has been estimated that at least one third of all organic compounds exhibit polymorphism.

In addition to polymorphic forms, compounds may occur in noncrystalline or amorphous forms. The energy required for a molecule of drug to escape from a crystal is much greater than is required to escape from an amorphous powder. Therefore, the amorphous form of a compound is always more soluble than a corresponding crystal form.

Evaluation of crystal structure, polymorphism, and solvate form is an important preformulation activity. The changes in crystal characteristics can influence bioavailability and chemical and physical stability and can have important implications in dosage form process functions. For example, it can be a significant factor relating to tablet formation because of flow and compaction



FIGURE 4.3 Mastersizer 2000E particle size analyzer. (Courtesy of Malvern Instruments Ltd.)

behaviors, among others. Various techniques are used to determine crystal properties. The most widely used methods are hot stage microscopy, thermal analysis, infrared spectroscopy, and X-ray diffraction.

Solubility

An important physicochemical property of a drug substance is solubility, especially aqueous system solubility. A drug must possess some aqueous solubility for therapeutic efficacy. For a drug to enter the systemic circulation and exert a therapeutic effect, it must first be in solution. Relatively insoluble compounds often exhibit incomplete or erratic absorption. If the solubility of the drug substance is less than desirable, consideration must be given to improve its solubility. The methods to accomplish this depend on the chemical nature of the drug and the type of drug product under consideration. Chemical modification of the drug into salt or ester forms is frequently used to increase solubility.

A drug's solubility is usually determined by the equilibrium solubility method, by which an excess of the drug is placed in a solvent and shaken at a constant temperature over a long period until equilibrium is obtained. Chemical analysis of the drug content in solution is performed to determine degree of solubility.

Solubility and Particle Size

Although solubility is normally considered a physicochemical constant, small increases in solubility can be accomplished by particle size reduction as described in the Physical Pharmacy Capsule 4.5, Solubility and Particle Size.

Solubility and pH

Another technique, if the drug is to be formulated into a liquid product, is adjustment of the pH of the solvent to enhance solubility. However, for many drug substances pH adjustment is not an effective means of improving

PHYSICAL PHARMACY CAPSULE 4.5

Solubility and Particle Size

The particle size and surface area of a drug exposed to a medium can affect actual solubility within reason, for example, in the following relationship:

$$\log \frac{S}{S_0} = \frac{2\gamma V}{2.303 RT r}$$

where

S is the solubility of the small particles,

S_0 is the solubility of the large particles,

γ is the surface tension,

V is the molar volume,

R is the gas constant,

T is the absolute temperature, and

r is the radius of the small particles.

The equation can be used to estimate the decrease in particle size required to increase solubility. For example, a desired increase in solubility of 5% would require an increase in the S/S_0 ratio to 1.05; that is, the left term in the equation would become $\log 1.05$. If a powder has a surface tension of 125 dynes per centimeter, molar volume of 45 cm^3 , and temperature of 27°C , what is the particle size required to obtain the 5% increase in solubility?

$$\log 1.05 = \frac{(2)(125)(45)}{(2.303)(8.314 \times 10^7)(300)r}$$

$$r = 9.238 \times 10^{-6} \text{ cm or } 0.09238 \mu$$

A number of factors are involved in actual solubility enhancement, and this is only an introduction to the general effects of particle size reduction.

solubility. Weak acidic or basic drugs may require extremes in pH that are outside accepted physiologic limits or that may cause stability problems with formulation ingredients. Adjustment of pH usually has little effect on the solubility of substances other than electrolytes. In many cases, it is desirable to use cosolvents or other techniques such as complexation, micronization, or solid dispersion to improve

aqueous solubility. A review of pH is provided in Physical Pharmacy Capsule 4.6, Principles of pH. The effect of pH on solubility is illustrated in Physical Pharmacy Capsule 4.7, Solubility and pH.

In recent years, more and more physicochemical information on drugs is being made available to pharmacists in routinely used reference books. This type of information is important



PHYSICAL PHARMACY CAPSULE 4.6

Principles of pH

pH is a critical variable in pharmaceuticals, and a basic understanding of its principles and measurement are important. Let's begin with a definition of the term pH. The p comes from the word power. The H, of course, is the symbol for hydrogen. Together, the term pH means the hydrogen ion exponent.

The pH of a substance is a measure of its acidity, just as a degree is a measure of temperature. A specific pH value tells the exact acidity. Rather than stating general ideas, such as cherry syrup is acidic or the water is hot, a specific pH value gives the same relative point of reference, thus providing more exact communication. "The cherry juice has a pH of 3.5" or "the water is at 80°C" provides an exact common language.

pH is defined in terms of the hydrogen ion activity:



$$\text{pH} = -\log_{10} a_{\text{H}^+} \text{ or } 10^{-\text{pH}} = a_{\text{H}^+}$$

pH equals the negative logarithm of the hydrogen ion activity, or the activity of the hydrogen ion is 10 raised to the exponent $-\text{pH}$. The latter expression renders the use of the p exponent more obvious. The activity is the effective concentration of the hydrogen ion in solution. The difference between effective and actual concentration decreases as one moves toward more dilute solutions, in which ionic interaction becomes progressively less important.

Normally, reference is made to the hydrogen ion when reference should be made to the hydronium ion (H_3O^+). It is a matter of convenience and brevity that only the hydrogen ion is mentioned, even though it is normally in its solvated form:



The complexing of the hydrogen ion by water affects activity and applies to other ions, which partially complex or establish an equilibrium with the hydrogen ion. In other words, equilibrium such as



complexes the hydrogen ion so that it is not sensed by the pH measuring system. This is why an acid-base titration is performed if the total concentration of acid (H^+) is needed. These effects on hydrogen ion activity are obvious, but other more subtle effects are involved in the correlation of activity and concentration.

The activity of the hydrogen ion can be defined by its relation to concentration (C_{H^+} , molality) and the activity coefficient f_{H^+} :

$$a_{\text{H}^+} = f_{\text{H}^+} \cdot C_{\text{H}^+}$$

If the activity coefficient is unity, activity is equal to concentration. This is nearly the case in dilute solutions, whose ionic strength is low. Since the objective of most pH measurements is to find a stable and reproducible reading that can be correlated to the results of some process, it is important to know what influences the activity coefficient and therefore the pH measurement.

PHYSICAL PHARMACY CAPSULE 4.6 CONT.

The factors that affect the activity coefficient are the temperature (T), the ionic strength (μ), the dielectric constant (ϵ), the ion charge (Z), the size of the ion in angstroms (\AA), and the density of the solvent (d). All of these factors are characteristics of the solution that relate the activity to the concentration by two main effects: the salt effect and the medium effect; the latter relates the influence that the solvent can have on the hydrogen ion activity. Thus, hydrogen activity is related to concentration through a salt effect and a solvent effect. Because of these influences, a sample pH value cannot be extrapolated to another temperature or dilution. If the pH value of a particular solution is known at 40°C , it is not automatically known at 25°C .

THE pH SCALE

In pure water, hydrogen and hydroxyl ion concentrations are equal at 10^{-7} M at 25°C . This is a neutral solution. Since most samples encountered have less than 1 M H^+ or OH^- , the extremes of pH 0 for acids and pH 14 for bases are established. Of course, with strong acids or bases, pH values below 0 and above 14 are possible but infrequently measured.

MEASUREMENT OF pH

The activity of the hydrogen ion in solution is measured with a glass electrode, a reference electrode, and a pH meter.

COMBINATION ELECTRODES

A combination electrode is a combination of the glass and reference electrodes into a single probe. The main advantage in using a combination electrode is with the measurement of small volume samples or samples in limited-access containers.

PHYSICAL PHARMACY CAPSULE 4.7

Solubility and pH

pH is one of the most important factors in the formulation process. Two areas of critical importance are the effects of pH on solubility and stability. The effect of pH on solubility is critical in the formulation of liquid dosage forms, from oral and topical solutions to intravenous solutions and admixtures.

The solubility of a weak acid or base is often pH dependent. The total quantity of a monoprotic weak acid (HA) in solution at a specific pH is the sum of the concentrations of both the free acid and salt (A^-) forms. If excess drug is present, the quantity of free acid in solution is maximized and constant because of its saturation solubility. As the pH of the solution increases, the quantity of drug in solution increases because the water-soluble ionizable salt is formed. The expression is



where K_a is the dissociation constant.

There may be a certain pH level reached where the total solubility (S_T) of the drug solution is saturated with respect to both the salt and acid forms of the drug, that is, the pH_{max} . The solution can be saturated with respect to the salt at pH values higher than this, but not with respect to the acid. Also, at pH values

PHYSICAL PHARMACY CAPSULE 4.7 CONT.

less than this, the solution can be saturated with respect to the acid but not to the salt. This is illustrated in the accompanying figure.

To calculate the total quantity of drug that can be maintained in solution at a selected pH, either of two equations can be used, depending on whether the product is to be in a pH region above or below the pH_{max} . The following equation is used when below the pH_{max} :

$$S_T = S_a \left(1 + \frac{K_a}{[H^+]} \right) \quad (\text{Equation 1})$$

The next equation is used when above the pH_{max} :

$$S_T = S'_a \left(1 + \frac{K_a}{[H^+]} \right) \quad (\text{Equation 2})$$

where

S_a is the saturation solubility of the free acid and

S'_a is the saturation solubility of the salt form.

EXAMPLE

A pharmacist prepares a 3.0% solution of an antibiotic as an ophthalmic solution and dispenses it to a patient. A few days later the patient returns the eye drops to the pharmacist because the product contains a precipitate. The pharmacist, checking the pH of the solution and finding it to be 6.0, reasons that the problem may be pH related. The physicochemical information of interest on the antibiotic includes the following:

Molecular weight	285 (salt) 263 (free acid)
3.0% solution of the drug	0.1053 M solution
Acid form solubility (S_a)	3.1 mg/mL (0.0118 M)
K_a	5.86×10^{-6}

Using Equation 1, the pharmacist calculates the quantity of the antibiotic in solution at a pH of 6.0 (*Note:* pH of 6.0 = $[H^+]$ of 1×10^{-6})


$$S_T = 0.0118[1+] = 0.0809 \text{ molar}$$

From this the pharmacist knows that at a pH of 6.0, a 0.0809-M solution can be prepared. However, the concentration that was to be prepared was a 0.1053-M solution; consequently, the drug will not be in solution at that pH. The pH may have been all right initially but shifted to a lower pH over time, resulting in precipitation of the drug. The question is at what pH (hydrogen ion concentration) the drug will remain in solution. This can be calculated using the same equation and information. The S_T value is 0.1053 M.

$$0.1053 = 0.0118 \left[1 + \frac{5.86 \times 10^{-6}}{[H^+]} \right]$$

$$[H^+] = 7.333 \times 10^{-7}, \text{ or a pH of } 6.135$$

The pharmacist prepares a solution of the antibiotic, adjusting the pH to above about 6.2, using a suitable buffer system, and dispenses the solution to the patient—with positive results.



PHYSICAL PHARMACY CAPSULE 4.7 CONT.

An interesting phenomenon concerns the close relationship of pH to solubility. At a pH of 6.0, only a 0.0809-M solution could be prepared, but at a pH of 6.13 a 0.1053-M solution could be prepared. In other words, a difference of 0.13 pH units resulted in

$$\frac{0.1053 - 0.0809}{0.0809} = 30.1\%$$

more drug going into solution at the higher pH than at the lower pH. In other words, a very small change in pH resulted in about 30% more drug going into solution. According to the figure, the slope of the curve would be very steep for this example drug, and a small change in pH (x-axis) results in a large change in solubility (y-axis). From this, it can be reasoned that if one observes the pH-solubility profile of a drug, it is possible to predict the magnitude of the pH change on its solubility.

In recent years, more and more physicochemical information on drugs is being made available to pharmacists in routinely used reference books. This type of information is important for pharmacists in different types of practice, especially those who compound and do pharmacokinetic monitoring.

for pharmacists in different types of practice, especially those involved in compounding and pharmacokinetic monitoring.

Dissolution

Variations in the biologic activity of a drug substance may be brought about by the rate at which it becomes available to the organism. In many instances, dissolution rate, or the time it takes for the drug to dissolve in the fluids at the absorption site, is the rate-limiting step in absorption. This is true for drugs administered orally in solid forms such as tablets, capsules, or suspensions, and for those administered intramuscularly. When the dissolution rate is the rate-limiting step, anything that affects it will also affect absorption. Consequently, dissolution rate can affect the onset, intensity, and duration of response and control the overall bioavailability of the drug from the dosage form, as discussed in the previous chapter.

The dissolution rate of drugs may be increased by decreasing the drug's particle size. It may also be increased by increasing its solubility in the diffusion layer. The most effective means of obtaining higher dissolution rates is to use a highly water-soluble salt of the parent substance. Although a soluble salt of a weak acid will precipitate as the free acid in the bulk phase of an acidic solution, such as gastric fluid, it will do so

in the form of fine particles with a large surface area.

The dissolution rates of chemical compounds are determined by two methods: the constant-surface method, which provides the intrinsic dissolution rate of the agent, and particulate dissolution, in which a suspension of the agent is added to a fixed amount of solvent without exact control of surface area.

The constant-surface method uses a compressed disc of known area. This method eliminates surface area and surface electrical charges as dissolution variables. The dissolution rate obtained by this method, the *intrinsic dissolution rate*, is characteristic of each solid compound and a given solvent in the fixed experimental conditions. The value is expressed as milligrams dissolved per minute per centimeters squared. It has been suggested that this value is useful in predicting probable absorption problems due to dissolution rate. In particulate dissolution, a weighed amount of powdered sample is added to the dissolution medium in a constant agitation system. This method is frequently used to study the influence of particle size, surface area, and excipients upon the active agent. Occasionally, the surface properties of the drug produce an inverse relationship of particle size to dissolution. In these instances, surface charge and/or agglomeration results in the reduced particle size form of the drug presenting

a lower effective surface area to the solvent due to incomplete wetting or agglomeration. Fick's laws describe the relationship of diffusion and dissolution of the active drug in the dosage form and when administered in the body, as shown in Physical Pharmacy Capsule 4.8, Fick's Laws of Diffusion and the Noyes–Whitney Equation.

Early formulation studies should include the effects of pharmaceutical ingredients on the dissolution characteristics of the drug substance.

Membrane Permeability

Modern preformulation studies include an early assessment of passage of drug molecules across biologic membranes. To produce a biologic response, the drug molecule must first cross a biologic membrane. The biologic membrane acts as a lipid barrier to most drugs and permits the absorption of lipid-soluble substances by passive diffusion, while lipid-insoluble



PHYSICAL PHARMACY CAPSULE 4.8

Fick's Laws of Diffusion and the Noyes-Whitney Equation

All drugs must diffuse through various barriers when administered to the body. For example, some drugs must diffuse through the skin, gastric mucosa, or some other barrier to gain access to the interior of the body. Parenteral drugs must diffuse through muscle, connective tissue, and so on, to get to the site of action; even intravenous drugs must diffuse from the blood to the site of action. Drugs must also diffuse through various barriers for metabolism and excretion.

Considering all the diffusion processes that occur in the body (passive, active, and facilitated), it is not surprising that the laws governing diffusion are important to drug delivery systems. In fact, diffusion is important not only in the body but also in some quality control procedures used to determine batch-to-batch uniformity of products (dissolution test for tablets based on the Noyes-Whitney equation, which can be derived from Fick's law).

When individual molecules move within a substance, diffusion is said to occur. This may occur as the result of a concentration gradient or by random molecular motion.

Probably the most widely used laws of diffusion are known as Fick's first and second laws. Fick's first law involving steady-state diffusion (where dc/dx does not change) is derived from the following expression for the quantity of material (M) flowing through a cross section of a barrier (S) in unit time (t) expressed as the flux (J):

$$J = dM/(Sdt)$$

Under a concentration gradient (dc/dx), Fick's first law can be expressed thus:

$$J = D[(C_1 - C_2)/h] \text{ or } J = -D(dc/dx)$$

where

- J is the flux of a component across a plane of unit area,
- C_1 and C_2 are the concentrations in the donor and receptor compartments,
- h is the membrane thickness, and
- D is the diffusion coefficient (or diffusivity).

The sign is negative, denoting that the flux is in the direction of decreasing concentration. The units of J are grams per square centimeter; C , grams per cubic centimeter; M , grams or moles; S , square centimeters; x , centimeters; and D , square centimeters per second.

D is appropriately called a diffusion coefficient, not a diffusion constant, as it is subject to change. D may change in value with increased concentrations. Also, D can be affected by temperature, pressure, solvent properties, and the chemical nature of the drug itself. To study the rate of change of the drug in the system, one needs an expression that relates the change in concentration with time at a definite location in place of the mass of drug diffusing across a unit area of barrier in unit time; this expression is known as Fick's second law. This law can be summarized as stating that the change in concentration in a particular place with time is proportional to the change in concentration gradient at that particular place in the system.

PHYSICAL PHARMACY CAPSULE 4.8 CONT.

In summary, Fick's first law relates to a steady-state flow, whereas Fick's second law relates to a change in concentration of drug with time, at any distance, or an unsteady state of flow.

The diffusion coefficients ($D \times 10^{-6}$) of various compounds in water (25°C) and other media have been determined as follows: ethanol, 12.5 cm^2 per second; glycine, 10.6 cm^2 per second; sodium lauryl sulfate, 6.2 cm^2 per second; glucose, 6.8 cm^2 per second.

The concentration of drug in the membrane can be calculated using the partition coefficient (K) and the concentration in the donor and receptor compartments.

$$K = (C_1/C_d) = (C_2/C_r)$$

where

C_1 and C_d are the concentrations in the donor compartment (g/cm^3)

and C_2 and C_r are the concentrations in the receptor compartment (g/cm^3).

K is the partition coefficient of the drug between the solution and the membrane. It can be estimated using the oil solubility of the drug versus the water solubility of the drug. Usually, the higher the partition coefficient, the more the drug will be soluble in a lipophilic substance. We can now write the expression:

$$dM/dt = [DSK(C_d - C_r)]/h$$

or in sink conditions,

$$dM/dt = DSKC_d/h = PSC_d$$

The permeability coefficient (centimeters per second) can be obtained by rearranging to:

$$P = DK/h$$

EXAMPLE 1

A drug passing through a 1-mm-thick membrane has a diffusion coefficient of $4.23 \times 10^{-7}\text{ cm}^2$ per second and an oil-water partition coefficient of 2.03. The radius of the area exposed to the solution is 2 cm, and the concentration of the drug in the donor compartment is 0.5 mg/mL. Calculate the permeability and the diffusion rate of the drug.

$$h = 1\text{ mm} = 0.1\text{ cm}$$

$$D = 4.23 \times 10^{-7}\text{ cm}^2/\text{second}$$

$$K = 2.03$$

$$r = 2\text{ cm}, S = \pi(2\text{ cm})^2 = 12.57\text{ cm}^2$$

$$C_d = 0.5\text{ mg/mL}$$

$$P = [(4.23 \times 10^{-7}\text{ cm}^2/\text{second})(2.03)]/0.1\text{ cm} = 8.59 \times 10^{-6}\text{ cm}/\text{second}$$

$$dM/dt = (8.59 \times 10^{-6}\text{ cm}/\text{second})(12.57\text{ cm}^2)(0.5\text{ mg/mL}) = 5.40 \times 10^{-5}\text{ mg}/\text{second}$$

$$(5.40 \times 10^{-5}\text{ mg}/\text{second})(3600\text{ second}/\text{hour}) = 0.19\text{ mg}/\text{hour}$$

In the dissolution of particles of drug, the dissolved molecules diffuse away from the individual particle body. An expression to describe this, derived from Fick's equations, is known as the Noyes and Whitney expression, proposed in 1897. It can be written as follows:

$$dC/dt = (DS/Vh)(C_s - C)$$

where

C is the concentration of drug dissolved at time t ,

D is the diffusion coefficient of the solute in solution,

S is the surface area of the exposed solid,

PHYSICAL PHARMACY CAPSULE 4.8 CONT.

V is the volume of solution,
 h is the thickness of the diffusion layer,
 C_s is the saturation solubility of the drug, and
 C is the concentration of solute in the bulk phase at a specific time, t.

It is common practice to use sink conditions in which C does not exceed about 20% of the solubility of the drug being investigated. Under these conditions, the expression simplifies to

$$dC/dt = (DSC_s/Vh)$$

and incorporating the volume of solution (V), the thickness of the diffusion layer (h), and the diffusivity coefficient (D) into a coefficient k (to take into account the various factors in the system), the expression becomes

$$dC/dt = kSC_s$$

As the factors are held constant, it becomes apparent that the dissolution rate of a drug can be proportional to the surface area exposed to the dissolution medium. A number of other expressions have been derived for specific application to various situations and conditions.

These relationships expressed as Fick's first and second laws and the Noyes-Whitney equation have great importance and relevance in pharmaceutical systems.

EXAMPLE 2

The following information was obtained using the USP 32-NF 27 dissolution apparatus I. The drug is soluble at 1 g in 3 mL of water, so sink conditions were maintained; the surface area of the tablet exposed was 1.5 cm² (obtained by placing the tablet in a special holder exposing only one side to the dissolution medium); and the dosage form studied was a 16-mg sustained-release tablet; the release pattern should be zero order. What is the rate of release of drug?

TIME (HOURS)	DRUG CONCENTRATION (mg/900 mL OF SOLUTION)	GRAPH OF RELEASE PROFILE
0.0	0.0	
0.5	1.0	
1.0	1.9	
2.0	4.1	
4.0	8.0	
6.0	11.8	
8.0	15.9	

In this problem, since the surface area (S) was maintained constant at 1.5 cm² and the solubility (C_s) of the drug is constant at 1 g in 3 mL of water, the plot of concentration versus time (t) yields a slope with a value of kSC_s, or k₂, expressing the rate of release of the drug as

$$dC/dt = kSC_s$$

The slope of the line = $\Delta y/\Delta x = (y_2 - y_1)/(x_2 - x_1)$
 $= (15.9 \text{ mg} - 0 \text{ mg})/(8.0 \text{ h} - 0 \text{ h})$
 $= 15.9/8 = 1.99 \text{ mg/h}$

Therefore, the rate of release of the sustained-release preparation is 1.99, or approximately 2 mg per hour. From this, the quantity of drug released at any time (t) can be calculated.

substances can diffuse across the barrier only with considerable difficulty if at all. The interrelationship of the dissociation constant, lipid solubility, and pH at the absorption site with the absorption characteristics of various drugs are the basis of the pH partition theory.

Data obtained from the basic physicochemical studies, specifically, pKa, solubility, and dissolution rate, provide an indication of absorption. To enhance these data, a technique using the everted intestinal sac may be used to evaluate absorption characteristics of drug substances. In this method, a piece of intestine is removed from an intact animal, is everted, and is filled with a solution of the drug substance, and the degree and rate of passage of the drug through the membrane sac are determined. This method allows evaluation of both passive and active transport.

In the latter stages of preformulation testing or early formulation studies, animals and humans must be studied to assess the absorption efficiency and pharmacokinetic parameters and to establish possible *in vitro* and *in vivo* correlation for dissolution and bioavailability.

Partition Coefficient

The use of the partition coefficient is described in some detail in Physical Pharmacy Capsule 4.9, Partition Coefficient. Inherent in this procedure is the selection of appropriate extraction solvents, drug stability, use of salting-out additives, and environmental concerns. The octanol–water partition coefficient is commonly used in formulation development. Following the illustrations provided earlier, it is defined as

$$P = \frac{(\text{Conc. of drug in octanol})}{(\text{Conc. of drug in water})}$$

P depends on the drug concentration only if the drug molecules have a tendency to associate in solution. For an ionizable drug, the following equation is applicable:

$$P = \frac{(\text{Conc. of drug in octanol})}{[1 - \alpha](\text{Conc. of drug in water})}$$

where α equals the degree of ionization.



PHYSICAL PHARMACY CAPSULE 4.9

Partition Coefficient

The oil–water partition coefficient is a measure of a molecule's lipophilic character; that is, its preference for the hydrophilic or lipophilic phase. If a solute is added to a mixture of two immiscible liquids, it will distribute between the two phases and reach an equilibrium at a constant temperature. The distribution of the solute (unaggregated and undissociated) between the two immiscible layers can be described thus:

$$K = C_U/C_L$$

where

- K is the distribution constant or partition constant,
- C_U is the concentration of the drug in the upper phase, and
- C_L is the concentration of the drug in the lower phase.

This information can be effectively used in the

1. Extraction of crude drugs
2. Recovery of antibiotics from fermentation broth
3. Recovery of biotechnology-derived drugs from bacterial cultures
4. Extraction of drugs from biologic fluids for therapeutic drug monitoring
5. Absorption of drugs from dosage forms (ointments, suppositories, transdermal patches)
6. Study of the distribution of flavoring oil between oil and water phases of emulsions
7. In other applications

PHYSICAL PHARMACY CAPSULE 4.9 CONT.

This basic relationship can be used to calculate the quantity of drug extracted from or remaining behind in a given layer and to calculate the number of extractions required to remove a drug from a mixture.

The concentration of drug found in the upper layer (U) of two immiscible layers is given thus:

$$U = Kr/(Kr + 1)$$

where

K is the distribution partition constant and

r is V_u/V_l , or the ratio of the volume of upper and lower phases.

The concentration of drug remaining in the lower layer (L) is given thus:

$$L = 1/(Kr + 1)$$

If the lower phase is successively extracted again with n equal volumes of the upper layer, each upper (U_n) contains the following fraction of the drug:

$$U_n = Kr/(Kr + 1)^n$$

where

U_n is the fraction contained in the n th extraction and

n is the n th successive volume.

The fraction of solute remaining in the lower layer (L_n) is given thus:

$$L_n = 1/(Kr + 1)^n$$

More efficient extractions are obtained using successive small volumes of the extraction solvent than single larger volumes. This can be calculated as follows when the same volume of extracting solvent is used in divided portions. For example, the fraction L_n remaining after the n th extraction:

$$L_n = \frac{1}{\left(\frac{Kr}{n} + 1\right)^n}$$

EXAMPLE 1

At 25°C and pH 6.8, the K for a second generation cephalosporin is 0.7 between equal volumes of butanol and the fermentation broth. Calculate the U, L, and L_n (using the same volume divided into fourths).

$$U = 0.7/(0.7 + 1) = 0.41, \text{ the fraction of drug extracted into the upper layer}$$

$$L = 1/(0.7 + 1) = 0.59, \text{ the fraction of drug remaining in the lower layer}$$

The total of the fractions in the U and L = 0.41 + 0.59 = 1.

If the fermentation broth is extracted with four successive extractions accomplished by dividing the quantity of butanol used into fourths, the quantity of drug remaining after the fourth extraction is

$$L_{4\text{th}} = \frac{1}{\left(\frac{0.7 \times 1}{4} + 1\right)^4} = 0.525$$

From this, the quantity remaining after a single volume, single extraction is 0.59, but when the single volume is divided into fourths and four successive extractions are done, the quantity remaining is 0.525; therefore, more was extracted using divided portions of the extracting solvent. Inherent in this procedure is the selection of appropriate extraction solvents, drug stability, use of salting-out additives, and environmental concerns.

pKa/Dissociation Constants

Among the physicochemical characteristics of interest is the extent of dissociation or ionization of drug substances. This is important because the extent of ionization has an important effect on the formulation and pharmacokinetic parameters of the drug. The extent of dissociation or ionization in many cases is highly dependent on the pH of the medium containing the drug. In formulation, often the vehicle is adjusted to a certain pH to obtain a certain level of ionization of the drug for solubility and stability. In the pharmacokinetic area, the extent of ionization of a drug has a strong effect on its extent of absorption,

distribution, and elimination. The dissociation constant, or pKa, is usually determined by potentiometric titration. For the practicing pharmacist, it is important in predicting precipitation in admixtures and in calculating the solubility of drugs at certain pH values. Physical Pharmacy Capsule 4.10, pKa/Dissociation Constants, presents a brief summary of dissociation and ionization concepts.

DRUG AND DRUG PRODUCT STABILITY


One of the most important activities of preformulation work is evaluation of the physical and chemical stability of the pure drug substance. It



PHYSICAL PHARMACY CAPSULE 4.10

pKa/Dissociation Constants

The dissociation of a weak acid in water is given by this expression:



At equilibrium, the reaction rate constants K_1 and K_2 are equal. This can be rearranged, and the dissociation constant defined as

$$K_a = \frac{K_1}{K_2} = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

where K_a is the acid dissociation constant.


For the dissociation of a weak base that does not contain a hydroxyl group, the following relationship can be used:



The dissociation constant is described by

$$K_a = \frac{[\text{H}^+][\text{B}]}{[\text{BH}^+]}$$

The dissociation of a hydroxyl-containing weak base,



The dissociation constant is described by

$$K_b = \frac{[\text{OH}^-][\text{BH}^+]}{[\text{B}]}$$

The hydrogen ion concentrations can be calculated for the solution of a weak acid using

$$[\text{H}^+] = \sqrt{K_a C}$$

Similarly, the hydroxyl ion concentration for a solution of a weak base is approximated by

$$[\text{OH}^-] = \sqrt{K_b C}$$

PHYSICAL PHARMACY CAPSULE 4.10 CONT.

Some practical applications of these equations are as follows.

EXAMPLE 1

The K_a of lactic acid is 1.387×10^{-4} at 25°C . What is the hydrogen ion concentration of a 0.02 M solution?

$$[\text{H}^+] = \sqrt{1.387 \times 10^{-4} \times 0.02} = 1.665 \times 10^{-3} \text{ G-ion/L}$$

EXAMPLE 2

The K_b of morphine is 7.4×10^{-7} . What is the hydroxyl ion concentration of a 0.02 M solution?

$$[\text{OH}^-] = \sqrt{7.4 \times 10^{-7} \times 0.02} = 1.216 \times 10^{-4} \text{ G-ion/L}$$

is essential that these initial studies be conducted using drug samples of known purity. The presence of impurities can lead to erroneous conclusions in such evaluations. Stability studies conducted in the preformulation phase include solid-state stability of the drug alone, solution phase stability, and stability in the presence of expected excipients. Initial investigation begins with knowledge of the drug's chemical structure, which allows the preformulation scientist to anticipate the possible degradation reactions.

Drug Stability: Mechanisms of Degradation

Chemical instability of medicinal agents may take many forms because the drugs in use today are of such diverse chemical constitution. Chemically, drug substances are alcohols, phenols, aldehydes, ketones, esters, ethers, acids, salts, alkaloids, glycosides, and others, each with reactive chemical groups having different susceptibilities to chemical instability. Chemically, the most frequently encountered destructive processes are hydrolysis and oxidation.

Hydrolysis is a solvolysis process in which (drug) molecules interact with water molecules to yield breakdown products. For example, aspirin, or acetylsalicylic acid, combines with a water molecule and hydrolyzes into one molecule of salicylic acid and one molecule of acetic acid.

Hydrolysis is probably the most important single cause of drug decomposition, mainly

because a great number of medicinal agents are esters or contain such other groupings as substituted amides, lactones, and lactams, which are susceptible to the hydrolytic process (2).

Another destructive process is oxidation, which destroys many drug types, including aldehydes, alcohols, phenols, sugars, alkaloids, and unsaturated fats and oils. Chemically, oxidation is loss of electrons from an atom or a molecule. Each electron lost is accepted by some other atom or molecule, reducing the recipient. In inorganic chemistry, oxidation is accompanied by an increase in the positive valence of an element, for example, ferrous (+2) oxidizing to ferric (+3). In organic chemistry, oxidation is frequently considered synonymous with the loss of hydrogen (dehydrogenation) from a molecule. Oxidation frequently involves free chemical radicals, which are molecules or atoms containing one or more unpaired electrons, such as molecular (atmospheric) oxygen ($\bullet\text{O}-\text{O}\bullet$) and free hydroxyl ($\bullet\text{OH}$). These radicals tend to take electrons from other chemicals, thereby oxidizing the donor.

Many of the oxidative changes in pharmaceutical preparations have the character of autoxidations. Autoxidations occur spontaneously under the initial influence of atmospheric oxygen and proceed slowly at first and then more rapidly. The process has been described as a type of chain reaction commencing with the union of oxygen with the drug molecule and continuing with a free radical of this oxidized molecule participating in the destruction of other drug molecules and so forth.

In drug product formulation work, steps are taken to reduce or prevent deterioration due to hydrolysis, oxidation, and other processes. These techniques are discussed later.

Drug and Drug Product Stability: Kinetics and Shelf Life

Stability is the extent to which a product retains within specified limits and throughout its period of storage and use (i.e., its shelf life) the same properties and characteristics that it possessed at the time of its manufacture.

Five types of stability concern pharmacists:

1. *Chemical*: Each active ingredient retains its chemical integrity and labeled potency within the specified limits.
2. *Physical*: The original physical properties, including appearance, palatability, uniformity, dissolution, and suspendability are retained.
3. *Microbiologic*: Sterility or resistance to microbial growth is retained according to the specified requirements. Antimicrobial agents retain effectiveness within specified limits.
4. *Therapeutic*: The therapeutic effect remains unchanged.
5. *Toxicologic*: No significant increase in toxicity occurs.

Chemical stability is important for selecting storage conditions (temperature, light, humidity), selecting the proper container for dispensing (glass vs. plastic, clear vs. amber or opaque, cap liners), and anticipating interactions when mixing drugs and dosage forms. Stability and expiration dating are based on reaction kinetics, that is, the study of the rate of chemical change and the way this rate is influenced by concentration of reactants, products, and other chemical species and by factors such as solvent, pressure, and temperature.

In considering chemical stability of a pharmaceutical, one must know the reaction order and reaction rate. The reaction order may be the overall order (the sum of the exponents of the concentration terms of the rate expression), or the order with respect to each reactant (the exponent of the individual concentration term in the rate expression).

Rate Reactions

The reaction rate is a description of the drug concentration with respect to time. Most commonly,

zero-order and first-order reactions are encountered in pharmacy. These are presented in Physical Pharmacy Capsule 4.11, Rate Reactions, along with some appropriate examples.

Q_{10} Method of Shelf Life Estimation

The Q_{10} method of shelf life estimation lets the pharmacist estimate shelf life for a product that has been stored or is going to be stored under a different set of conditions. It is explained in Physical Pharmacy Capsule 4.12, Q_{10} Method of Shelf Life Estimation.

Enhancing Stability of Drug Products

Many pharmaceutical ingredients may be used to prepare the desired dosage form of a drug substance. Some of these agents may be used to achieve the desired physical and chemical characteristics of the product or to enhance its appearance, odor, and taste. Other substances may be used to increase the stability of the drug substance, particularly against hydrolysis and oxidation. In each instance, the added pharmaceutical ingredient must be compatible with and must not detract from the stability of the drug substance.

There are several approaches to the stabilization of pharmaceutical preparations containing drugs subject to hydrolysis. Perhaps the most obvious is the reduction or elimination of water from the pharmaceutical system. Even solid dosage forms containing water-labile drugs must be protected from humidity in the atmosphere. This may be accomplished by applying a waterproof protective coating over tablets or by keeping the drug in a tightly closed container. It is fairly common to detect hydrolyzed aspirin by noticing an odor of acetic acid upon opening a bottle of aspirin tablets. In liquid preparations, water can frequently be replaced or reduced in the formulation through the use of substitute liquids such as glycerin, propylene glycol, and alcohol. In certain injectable products, anhydrous vegetable oils may be used as the drug's solvent to reduce the chance of hydrolytic decomposition.

Decomposition by hydrolysis may be prevented in other liquid drugs by suspending them in a nonaqueous vehicle rather than dissolving them in an aqueous solvent. In still other instances, particularly for certain unstable



PHYSICAL PHARMACY CAPSULE 4.11

Rate Reactions

ZERO-ORDER RATE REACTIONS

If the loss of drug is independent of the concentration of the reactants and constant with respect to time (i.e., 1 mg/mL/hour), the rate is called zero order. The mathematical expression is

$$\frac{-dC}{dt} = k_0$$

where k_0 is the zero-order rate constant [concentration(C)/time(t)].

The integrated and more useful form of the equation:

$$C = -k_0t + C_0$$

where C_0 is the initial concentration of the drug.

The units for a zero rate constant k_0 are concentration per unit time, such as moles per liter-second or milligrams per milliliter per minute.

It is meaningless to attempt to describe the time required for *all* material in a reaction to decompose, that is, infinity. Therefore, reaction rates are commonly described by k or by their half-life, $t_{1/2}$.

The half-life equation for a zero-order reaction:

$$t_{1/2} = (1/2)(C_0/k_0)$$

If the C_0 changes, the $t_{1/2}$ changes. There is an inverse relationship between the $t_{1/2}$ and k .

EXAMPLE 1

A drug suspension (125 mg/mL) decays by zero-order kinetics with a reaction rate constant of 0.5 mg/mL/hour. What is the concentration of intact drug remaining after 3 days (72 hours), and what is its $t_{1/2}$?

$$C = -(0.5 \text{ mg/mL/hour})(72 \text{ hour}) + 125 \text{ mg/mL}$$

$$C = 89 \text{ mg/mL after 3 days}$$

$$t_{1/2} = 1/2(125 \text{ mg/mL})/(0.5 \text{ mg/mL/hour})$$

$$t_{1/2} = 125 \text{ hours}$$

EXAMPLE 2

How long will it take for the suspension to reach 90% of its original concentration?

$$90\% \times 125 \text{ mg/mL} = 112.5 \text{ mg/mL}$$

$$t = \frac{C - C_0}{-k_0} = \frac{112.5 \text{ mg/mL} - 125 \text{ mg/mL}}{-0.5 \text{ mg/mL/hour}} = 25 \text{ hours}$$

Drug suspensions are examples of pharmaceuticals that ordinarily follow zero-order kinetics for degradation.

FIRST-ORDER RATE REACTIONS

If the loss of drug is directly proportional to the concentration remaining with respect to time, it is called a first-order reaction and has the units of reciprocal time, that is, time^{-1} . The mathematical expression is

$$\frac{-dC}{dt} = kC$$

where

C is the concentration of intact drug remaining,

t is time,

(dC/dt) is the rate at which the intact drug degrades, and

k is the specific reaction rate constant.

PHYSICAL PHARMACY CAPSULE 4.11 CONT.

The integrated and more useful form of the equation:

$$\log C = \frac{-kt}{2.303} + \log C_0$$

where C_0 is the initial concentration of the drug.

In natural log form, the equation is

$$\ln C = -kt + \ln C_0$$

The units of k for a first-order reaction are per unit of time, such as per second.

The half-life equation for a first-order reaction is

$$t_{1/2} = 0.693/k$$

and can be easily derived from the first-order equation by substituting values of $C = 50\%$ and $C_0 = 100\%$, representing a decrease in concentration by 50%.

EXAMPLE 3

An ophthalmic solution of a mydriatic drug at 5 mg/mL exhibits first-order degradation with a rate of 0.0005/day. How much drug will remain after 120 days, and what is its half-life?

$$\ln C = -(0.0005/\text{day})(120) + \ln(5 \text{ mg/mL})$$

$$\ln C = -0.06 + 1.609$$

$$\ln C = 1.549$$

$$C = 4.71 \text{ mg/mL}$$

$$t_{1/2} = 0.693/0.0005/\text{day}$$

$$t_{1/2} = 1,386 \text{ days}$$

EXAMPLE 4

In Example 3, how long will it take for the drug to degrade to 90% of its original concentration?

$$90\% \text{ of } 5 \text{ mg/mL} = 4.5 \text{ mg/mL}$$

$$\ln 4.5 \text{ mg/mL} = -(0.0005/\text{day})t + \ln(5 \text{ mg/mL})$$

$$t = \frac{\ln 4.5 \text{ mg/mL} - \ln 5 \text{ mg/mL}}{-0.0005/\text{day}}$$

$$t = 210 \text{ days}$$

ENERGY OF ACTIVATION: ARRHENIUS EQUATION

Stability projections for shelf life (t_{90} , or the time required for 10% of the drug to degrade with 90% of the intact drug remaining) are commonly based on the Arrhenius equation:

$$\log \frac{k_2}{k_1} = \frac{E_a(T_2 - T_1)}{2.3RT_1T_2}$$

which relates the reaction rate constants (k) to temperatures (T) with the gas constant (R) and the energy of activation (E_a).

The relationship of the reaction rate constants at two different temperatures provides the energy of activation for the degradation. By performing the reactions at elevated temperatures instead of allowing the process to proceed slowly at room temperature, the E_a can be calculated and a k value for room temperature determined with the Arrhenius equation.

PHYSICAL PHARMACY CAPSULE 4.11 CONT.

EXAMPLE 5

The degradation of a new cancer drug follows first-order kinetics and has first-order degradation rate constants of 0.0001 per hour at 60°C and 0.0009 at 80°C. What is its E_a ?

$$\log = \frac{(0.0009)}{(0.0001)} = \frac{E_a(353 - 333)}{(2.3)(1.987)(353)(333)}$$

$$E_a = 25,651 \text{ kcal/mol}$$



PHYSICAL PHARMACY CAPSULE 4.12

Q₁₀ Method of Shelf Life Estimation

The Q_{10} approach, based on E_a , which is independent of reaction order, is described as

$$Q_{10} = e^{\{(E_a/R)[(1/T + 10) - (1/T)]\}}$$

where

E_a is the energy of activation,
 R is the gas constant, and
 T is the absolute temperature.

In usable terms, Q_{10} , the ratio of two different reaction rate constants, is defined thus:

$$Q_{10} = \frac{K_{(T+10)}}{K_T}$$

The commonly used Q values of 2, 3, and 4 relate to the energies of activations of the reactions for temperatures around room temperature (25°C). For example, a Q value of 2 corresponds to an E_a (kcal/mol) of 12.2, a Q value of 3 corresponds to an E_a of 19.4, and a Q value of 4 corresponds to an E_a of 24.5. Reasonable estimates can often be made using the value of 3.

The equation for Q_{10} shelf life estimates is

$$t_{90}(T_2) = \frac{t_{90}(T_1)}{Q_{10}^{(\Delta T/10)}}$$

where

$t_{90} T_2$ is the estimated shelf life,
 $t_{90} T_1$ is the given shelf life at a given temperature, and
 ΔT is the difference in the temperatures T_1 and T_2 .

As is evident from this relationship, an increase in ΔT will decrease the shelf life and a decrease in ΔT will increase shelf life. This is the same as saying that storing at a warmer temperature will shorten the life of the drug and storing at a cooler temperature will increase the life of the drug.

EXAMPLE 1

An antibiotic solution has a shelf life of 48 hours in the refrigerator (5°C). What is its estimated shelf life at room temperature (25°C)?

Using a Q value of 3, we set up the relationship as follows:

$$t_{90}(T_2) = \frac{t_{90}(T_1)}{Q_{10}^{(\Delta T/10)}} = \frac{48}{3^{[(25-5)/10]}} = \frac{48}{3^2} = 5.33 \text{ hours}$$

PHYSICAL PHARMACY CAPSULE 4.12 CONT.

EXAMPLE 2

An ophthalmic solution has a shelf life of 6 hours at room temperature (25°C). What is the estimated shelf life in a refrigerator at 5°C? (*Note:* Since the temperature is decreasing, ΔT will be negative.)

$$t_{90}(T_2) = \frac{6}{3^{[(5-25)/10]}} = \frac{6}{3^{-2}} = 6 \times 3^2 = 54 \text{ hours}$$

These are estimates, and actual energies of activation can often be obtained from the literature for more exact calculations.

antibiotic drugs, when an aqueous preparation is desired, the drug may be supplied to the pharmacist in a dry form for *reconstitution* by adding a specified volume of purified water just before dispensing. The dry powder is actually a mixture of the antibiotic, suspending agents, flavorants, and colorants; when reconstituted by the pharmacist, it remains stable for the period over which the preparation is normally consumed. Refrigeration is advisable for most preparations considered subject to hydrolysis. Together with temperature, pH is a major determinant of the stability of a drug prone to hydrolytic decomposition. Hydrolysis of most drugs depends on the relative concentrations of the hydroxyl and hydronium ions, and a pH at which each drug is optimally stable can be easily determined. For most hydrolyzable drugs, optimum stability is on the acid side, somewhere between pH 5 and 6. Therefore, through judicious use of buffering agents, the stability of otherwise unstable compounds can be increased. Buffers are used to maintain a certain pH, as described in Physical Pharmacy Capsule 4.13, Buffer Capacity.

Pharmaceutically, oxidation of a susceptible drug substance is most likely to occur when it is not kept dry in the presence of oxygen or when it is exposed to light or combined with other chemical agents without proper regard to their influence on oxidation. Oxidation of a chemical in a pharmaceutical preparation is usually accompanied by an alteration in the color of that preparation. It may also result in precipitation or a change in odor.

The oxidative process is diverted and the stability of the drug is preserved by agents called *antioxidants*, which react with one or more compounds in the drug to prevent progress of the chain reaction. In general, antioxidants act by

providing electrons and easily available hydrogen atoms that are accepted more readily by the free radicals than are those of the drug being protected. Various antioxidants are employed in pharmacy. Among those most frequently used in aqueous preparations are sodium sulfite (Na_2SO_3 , at high pH values), sodium bisulfite (NaHSO_3 , at intermediate pH values), sodium metabisulfite ($\text{Na}_2\text{S}_2\text{O}_5$ at low pH values), hypophosphorous acid (H_3PO_2), and ascorbic acid. In oleaginous (oily or unctuous) preparations, alpha-tocopherol, butyl hydroxy anisole, and ascorbyl palmitate find application.

In June 1987, U.S. Food and Drug Administration (FDA) labeling regulations went into effect requiring a warning about possible allergic-type reactions, including anaphylaxis, in the package insert for prescription drugs to whose final dosage form sulfites have been added. Sulfites are used as preservatives in many injectable drugs, such as antibiotics and local anesthetics. Some inhalants and ophthalmic preparations also contain sulfites, but relatively few oral drugs contain these chemicals. The purpose of the regulation is to protect the estimated 0.2% of the population who are subject to allergic reactions to the chemicals. Many sulfite-sensitive persons have asthma or other allergic conditions. Previous to the regulations dealing with prescription medication, the FDA issued regulations for the use of sulfites in food. Asthmatics and other patients who may be sulfite sensitive should be reminded to read the labels of packaged foods and medications to check for the presence of these agents. Sulfite agents covered by the regulations are potassium bisulfite, potassium metabisulfite, sodium bisulfite, sodium metabisulfite, sodium sulfite, and sulfur dioxide. The FDA permits the use of

PHYSICAL PHARMACY CAPSULE 4.13

Buffer Capacity

pH, buffers, and buffer capacity are especially important in drug product formulation, since they affect the drug's solubility, activity, absorption, and stability and the patient's comfort.

A buffer is a system, usually an aqueous solution, that can resist changes in pH upon addition of an acid or base. Buffers are composed of a weak acid and its conjugate base or a weak base and its conjugate acid. Buffers are prepared by one of these processes:

1. Mixing a weak acid and its conjugate base or a weak base and its conjugate acid
2. Mixing a weak acid and a strong base to form the conjugate base or a weak base and a strong acid to form the conjugate acid

Using the Henderson-Hasselbach equation:

$$\text{pH} = \text{pK}_a + \log (\text{base/acid})$$

Remember that the acid is the proton donor and the base is the proton acceptor.

EXAMPLE 1

A buffer is prepared by mixing 100 mL of 0.2 M phosphoric acid with 200 mL of 0.08 M sodium phosphate monobasic. What is the pH of this buffer? (K_a of phosphoric acid = 7.5×10^{-3})

$$\text{Moles acid} = (0.2 \text{ mol}/1000 \text{ mL})(100 \text{ mL}) = 0.02 \text{ mol}; (0.02 \text{ mol})/(0.3 \text{ L}) = 0.067 \text{ M}$$

$$\text{Moles base} = (0.08 \text{ mol}/1000 \text{ mL})(200 \text{ mL}) = 0.016 \text{ mol}; (0.016 \text{ mol})/(0.3 \text{ L}) = 0.053 \text{ M}$$

$$\text{pK}_a = -\log 7.5 \times 10^{-3} = 2.125$$

$$\text{pH} = 2.125 + \log (0.016 \text{ mol}/0.02 \text{ mol}) = 2.028$$

EXAMPLE 2

Determine the pH of the buffer prepared as shown:

Sodium acetate 50 g

Conc. HCl 10 mL

Water q.s. 2 L

Helpful numbers:

$$\text{pK}_a \text{ acetic acid} = 4.76$$

$$\text{m.w. sodium acetate} = 82.08$$

$$\text{m.w. acetic acid} = 60.05$$

$$\text{m.w. HCl} = 36.45$$

$$\text{Conc. HCl, 44\% HCl w/v}$$



$$(0.609 \text{ mol}) \quad (0.121 \text{ mol}) \quad (0.121 \text{ mol}) \quad (0.121 \text{ mol}) \quad (0.488 \text{ mol})$$

$$\text{HCl: } \{(10 \text{ mL}) [(44 \text{ g})/(100 \text{ mL})] (1 \text{ mol})/(36.45 \text{ g})\} = 0.121 \text{ mol}$$

$$\text{NaAc: } \{(50 \text{ g}) [(1 \text{ mol})/(82.08 \text{ g})]\} = 0.609 \text{ mol} \quad (0.609 \text{ mol}) - (0.121 \text{ mol}) = 0.488 \text{ mol}$$

$$\text{pH} = 4.76 + \log (0.488 \text{ mol})/(0.121 \text{ mol}) = 5.367$$

The ability of a buffer solution to resist changes in pH upon the addition of an acid or a base is called buffer capacity (β) and is defined thus:

$$\beta = \Delta B/\Delta \text{pH}$$

where

ΔB is molar concentration of acid or base added,

ΔpH is change in pH due to addition of acid or base, and

ΔpH can be determined experimentally or calculated using the Henderson-Hasselbach equation.

PHYSICAL PHARMACY CAPSULE 4.13 CONT.

EXAMPLE 3

If 0.2 mole of HCl is added to a 0.015 M solution of ammonium hydroxide and the pH falls from 9.5 to 8.9, what is the buffer capacity?

$$\begin{aligned}\Delta\text{pH} &= 9.5 - 8.9 = 0.6 \\ \Delta\text{B} &= 0.2 \text{ mol/L} = 0.2 \text{ M} \\ \beta &= 0.2 \text{ M}/0.6 = 0.33 \text{ M}\end{aligned}$$

EXAMPLE 4

If 0.002 mole of HCl is added to the buffer in Example 1, what is its buffer capacity? After adding 0.002 mole HCl:

$$\begin{aligned}\text{H}_3\text{PO}_4 &: 0.02 \text{ mol} + 0.002 \text{ mol} = 0.022 \text{ mol} \\ \text{NaH}_2\text{PO}_4 &: 0.016 \text{ mol} - 0.002 \text{ mol} = 0.014 \text{ mol} \\ \text{pH} &= 2.125 + \log(0.014 \text{ mol}/0.022 \text{ mol}) = 1.929 \\ \Delta\text{pH} &= 2.028 - 1.929 = 0.099 \\ \Delta\text{AB} &= 0.002 \text{ mol}/0.3 \text{ L} = 0.0067 \text{ M} \\ \beta &= 0.0067 \text{ M}/0.099 = 0.067 \text{ M}\end{aligned}$$

Another approach to calculating buffer capacity involves the use of Van Slyke's equation:

$$\beta = 2.3C \{ \text{Ka}[\text{H}^+]/(\text{Ka}[\text{H}^+])^2 \}$$

where

C is the sum of the molar concentrations of the acid and base, and $[\text{H}^+] = 10^{-\text{pH}}$.

EXAMPLE 5

What is the Van Slyke buffer capacity of the buffer prepared in Example 1?

$$\begin{aligned}C &= 0.0067 \text{ M} + 0.0053 \text{ M} = 0.12 \text{ M} \\ \text{Ka} &= 7.5 \times 10^{-3} \\ [\text{H}^+] &= 10^{-2.028} = 9.38 \times 10^{-3} \text{ M} \\ \beta &= 2.3(0.12 \text{ M}) \{ [(7.5 \times 10^{-3} \text{ M})(9.38 \times 10^{-3} \text{ M})] / [(7.5 \times 10^{-3} \text{ M}) / (9.38 \times 10^{-3} \text{ M})^2] \} = 0.68 \text{ M}\end{aligned}$$

sulfites in prescription products, with the proper labeling, because there are no generally suitable substitutes for sulfites to maintain potency in certain medications. Some but not all epinephrine injections contain sulfites.

The proper use of antioxidants permits their specific application only after appropriate biomedical and pharmaceutical studies. In certain instances, other pharmaceutical additives can inactivate a given antioxidant. In other cases, certain antioxidants can react chemically with the drugs they were intended to stabilize without a noticeable change in the appearance of the preparation.

Because oxygen may adversely affect their stability, certain pharmaceuticals require an oxygen-free atmosphere during preparation and storage. Oxygen may be present in pharmaceutical liquids in the airspace within the container or may be dissolved in the liquid vehicle. To avoid these exposures, oxygen-sensitive drugs may be prepared in the dry state and packaged in sealed containers with the air replaced by an inert gas such as nitrogen, as may liquid preparations. This is common practice in commercial production of vials and ampuls of easily oxidizable preparations intended for parenteral use.

TABLE 4.2 SOME USP DRUGS AND PREPARATIONS ESPECIALLY SUBJECT TO CHEMICAL OR PHYSICAL DETERIORATION


PREPARATION	CATEGORY	MONOGRAPH OR LABEL WARNING
Epinephrine bitartrate ophthalmic solution	Adrenergic	Do not use inhalation, injection, nasal, or ophthalmic solution if it is brown or contains a precipitate
Epinephrine inhalation solution		
Epinephrine injection		
Epinephrine nasal solution		
Epinephrine ophthalmic solution		
Isoproterenol sulfate inhalation, solution	Adrenergic (bronchodilator)	Do not use inhalation or injection if it is pink to brown or contains a precipitate
Isoproterenol inhalation solution		
Nitroglycerin tablets	Antianginal	To prevent loss of potency, keep in original container or supplemental container specifically labeled suitable for nitroglycerin tablets
Paraldehyde	Hypnotic	Subject to oxidation to form acetic acid

Trace metals originating in the drug, solvent, container, or stopper are a constant source of difficulty in preparing stable solutions of oxidizable drugs. The rate of formation of color in epinephrine solutions, for instance, is greatly increased by the presence of ferric, ferrous, cupric, and chromic ions. Great care must be taken to eliminate these trace metals from labile preparations by thorough purification of the source of the contaminant or by chemically complexing or binding the metal through the use of specialized agents that make it chemically unavailable for participation in the oxidative process. These chelating agents are exemplified by calcium disodium edetate and ethylenediaminetetraacetic acid.

Light can also act as a catalyst to oxidation reactions, transferring its energy (photons) to drug molecules, making the latter more reactive through increased energy capability. As a precaution against acceleration of oxidation, sensitive preparations are packaged in light-resistant or opaque containers.

Because most drug degradations proceed more rapidly as temperature increases, it is also advisable to maintain oxidizable drugs in a cool place. Another factor that can affect the stability of an oxidizable drug in solution is the pH of the preparation. Each drug must be maintained in solution at the pH most favorable to its stability. This varies from preparation to preparation and must be determined on an individual basis for the drug in question.

Statements in the United States Pharmacopeia (USP), as with those in Table 4.2, warn of the oxidative decomposition of drugs and preparations. In some instances, the specific agent to employ as a stabilizer is mentioned in the monograph, and in others the term “suitable stabilizer” is used. An example in which a particular agent is designated for use is in the monograph for potassium iodide oral solution, USP. Potassium iodide in solution is prone to photocatalyzed oxidation and the release of free iodine, with a resultant yellow-to-brown discoloration of the solution. The use of light-resistant containers is essential to its stability. As a further precaution against decomposition if the solution is not to be used within a short time, the USP recommends the addition of 0.5 mg of sodium thiosulfate for each gram of potassium iodide. In the event free iodine is released during storage, the sodium thiosulfate converts it to colorless and soluble sodium iodide:



In summary, for easily oxidizable drugs, the formulation pharmacist may stabilize the preparation by the selective exclusion from the system of oxygen, oxidizing agents, trace metals, light, heat, and other chemical catalysts to the oxidation process. Antioxidants, chelating agents, and buffering agents may be added to create and maintain a favorable pH.

In addition to oxidation and hydrolysis, destructive processes include polymerization,

chemical decarboxylation, and deamination. However, these processes occur less frequently and are peculiar to only small groups of chemical substances. Polymerization is a reaction between two or more identical molecules that forms a new and generally larger molecule. Formaldehyde is an example of a drug capable of polymerization. In solution it may polymerize to paraformaldehyde $(\text{CH}_2\text{O})_n$, a slowly soluble white crystalline substance that may cloud the solution. The formation of paraformaldehyde is enhanced by cool storage, especially in solutions with high concentrations of formaldehyde. The official formaldehyde solution contains approximately 37% formaldehyde and according to the USP, should be stored at temperatures not below 15°C (59°F). If the solution becomes cloudy upon standing in a cool place, it usually may be cleared by gentle warming. Formaldehyde is prepared by the limited oxidation of methanol (methyl alcohol), and the USP permits a residual amount of this material to remain in the final product, since it can retard the formation of paraformaldehyde. Formaldehyde solution must be maintained in a tight container because oxidation of the formaldehyde yields formic acid.



Other organic drug molecules may be degraded through processes in which one or more of their active chemical groups are removed. These processes may involve various catalysts, including light and enzymes. Decarboxylation and deamination are examples of such processes; the former is decomposition of an organic acid ($\text{R}\cdot\text{COOH}$) and release of carbon dioxide gas, and the latter is removal of the nitrogen-containing group from an organic amine. For example, insulin, a protein, deteriorates rapidly in acid solutions as a result of extensive deamination (3). Thus, most preparations of insulin are neutralized to reduce the rate of decomposition.

Stability Testing

FDA's Current Good Manufacturing Practice regulations include sections on stability and stability testing of pharmaceutical components and finished pharmaceutical products. In addition, FDA and International Conference on

Harmonization guidelines and guidances provide working recommendations to support the regulatory requirements. Among these are the following (4):

- “Stability Testing of New Drug Substances and Products”
- “Quality of Biotechnological Products: Stability Testing of Biotechnology/Biological Drug Products”
- “Photostability Testing of New Drug Substances and Products”
- “Stability Testing of New Dosage Forms”

Drug and drug product stability testing during every stage of development is critical to the quality of the product. Drug stability is important during preclinical testing and in clinical (human) trials to obtain a true and accurate assessment of the product being evaluated. For a marketed drug product, assurance of stability is vital to its safety and effectiveness during the course of its shelf life and use.

The FDA-required demonstration of drug stability is necessarily different for each stage of drug development, such as for a 2-week preclinical study, an early Phase I study, a limited Phase II trial, a pivotal Phase III clinical study, or for a new drug application. As a drug development program progresses, so do the requisite data to demonstrate and document the product's stability profile. Before approval for marketing, a product's stability must be assessed with regard to its formulation; the influence of its pharmaceutical ingredients; the influence of the container and closure; the manufacturing and processing conditions (e.g., heat); packaging components; conditions of storage; anticipated conditions of shipping, temperature, light, and humidity; and anticipated duration and conditions of pharmacy shelf life and patient use. Holding intermediate product components (such as drug granulations for tablets) for long periods before processing into finished pharmaceutical products can affect the stability of both the intermediate component and the finished product. Therefore, in-process stability testing, including retesting of intermediate components, is important.

Product containers, closures, and other packaging features must be considered in stability testing. For instance, tablets or capsules packaged in glass or plastic bottles require different stability test protocols from those for blister packs or strip packaging. Drugs particularly subject to

hydrolysis or oxidative decomposition must be evaluated accordingly. And sterile products must meet sterility test standards to ensure protection against microbial contamination. All preservatives must be tested for effectiveness in the finished product.

As noted elsewhere in this section, drug products must meet stability standards for long-term storage at room temperature and relative humidity. Products are also subjected to accelerated stability studies as an indication of shelf life stability. It is an FDA requirement that if the data are not submitted in the approved application, the first three postapproval production batches of a drug substance must be subjected to long-term stability studies and the first three postapproval production batches of drug product must be subjected to both long-term and accelerated stability studies (5, 6).

Drug instability in pharmaceutical formulations may be detected in some instances by a change in the physical appearance, color, odor, taste, or texture of the formulation, whereas in other instances chemical changes may not be self-evident and may be ascertained only through chemical analysis. Scientific data pertaining to the stability of a formulation can lead to prediction of the expected shelf life of the proposed product, and when necessary to redesign of the drug (e.g., into more stable salt or ester form) and to reformulation of the dosage form. Obviously, the rate at which a drug product degrades is of prime importance. The study of the rate of chemical change and the way it is influenced by such factors as the concentration of the drug or reactant, the solvent, temperature and pressure, and other chemical agents in the formulation is reaction kinetics.

In general, a kinetic study begins by measuring the concentration of the drug at given intervals under a specific set of conditions including temperature, pH, ionic strength, light intensity, and drug concentration. The measurement of the drug's concentration at the various times reveals the stability or instability of the drug under the specified conditions with the passage of time. From this starting point, each of the original conditions may be varied to determine the influence of such changes on the drug's stability. For example, the pH of the solution may be changed while the temperature, light intensity, and original drug concentration are held constant.

The findings may be presented graphically, by plotting the drug concentration as a function of time. From the experimental data, the reaction rate may be determined and a rate constant and half-life calculated.

The use of exaggerated conditions of temperature, humidity, light, and others to test the stability of drug formulations is termed accelerated stability testing. Accelerated temperature stability studies, for example, may be conducted for 6 months at 40°C with 75% relative humidity. If a significant change in the product occurs under these conditions, lesser temperature and humidity may be used, such as 30°C and 60% relative humidity. Short-term accelerated studies are used to determine the most stable of the proposed formulations for a drug product. In stress testing, temperature elevations in 10°C increments higher than used in accelerated studies are employed until chemical or physical degradation. Once the most stable formulation is ascertained, its long-term stability is predicted from the data generated from continuing stability studies. Depending on the types and severity of conditions employed, it is fairly common to maintain samples under exaggerated conditions of both temperature and varying humidity for 6 to 12 months. Such studies lead to the prediction of shelf life for a drug product.

In addition to the accelerated stability studies, drug products are subjected to long-term stability studies under the usual conditions of transport and storage expected during product distribution. In conducting these studies, the different national and international climate zones to which the product may be subjected must be borne in mind and expected variances in conditions of temperature and humidity included in the study design. Geographic regions are defined by zones: zone I, temperate; zone II, subtropical; zone III, hot and dry; and zone IV, hot and humid. A given drug product may encounter more than a single zone of temperature and humidity variations during its production and shelf life. Furthermore, it may be warehoused, transported, placed on a pharmacy's shelf, and subsequently in the patient's medicine cabinet, over a varying time course and at a wide range of temperature and humidity. In general, however, the long-term (12 months minimum) testing of new drug entities is conducted at 25°C ± 2°C and at a relative humidity of 60% ± 5%.

Samples maintained under these conditions may be retained for 5 years or longer, during which time they are observed for physical signs of deterioration and chemically assayed. These studies, considered with the accelerated stability studies previously performed, lead to a more precise determination of drug product stability, actual shelf life, and the possible extension of expiration dating.

When chemical degradation products are detected, the FDA requires the manufacturer to report their chemical identity, including structure, mechanism of formation, physical and chemical properties, procedures for isolation and purification, specifications and directions for determination at levels expected to be present in the pharmaceutical product, and their pharmacologic action and biologic significance, if any.

Physical Pharmacy Capsule 4.14 Analytical Methods and Standard Curves discusses some analytical methods and standard curve construction used in studies of this type.

In addition, signs of degradation of the specific dosage forms must be observed and reported. For the various dosage forms, this includes the following (1):

Tablets: Appearance (cracking, chipping, mottling), friability, hardness, color, odor, moisture content, clumping, disintegration, and dissolution.

Capsules: Moisture tackiness, color, appearance, shape, brittleness, and dissolution.

Oral solutions and suspensions: Appearance, precipitation, pH, color, odor, redispersibility (suspensions), and clarity (solutions).

Oral powders: Appearance, color, odor, and moisture.

Metered-dose inhalation aerosols: Delivered dose per actuation, number of metered doses, color, particle size distribution, loss of propellant, pressure, valve corrosion, spray pattern, and absence of pathogenic microorganisms.

Topical nonmetered aerosols: Appearance, odor, pressure, weight loss, net weight dispensed, delivery rate, and spray pattern.

Topical creams, ointments, lotions, solutions, and gels: Appearance, color, homogeneity, odor, pH, resuspend-



PHYSICAL PHARMACY CAPSULE 4.14

Analytical Methods and Standard Curves

Any study involving concentration of a drug requires an analytical method and the development of standard curves. There are numerous analytical methods used in pharmacy. It is important for pharmacists to have a basic understanding of pharmaceutical analysis to ensure that valid results are obtained when tests are being conducted. It is important to know (a) when to test, (b) what to test, (c) what method(s) to use, (d) how to interpret the results, (e) the limits of the test, and (f) the importance of analytical testing in the overall quality program in pharmacy.

The goal in analytical testing is to produce results as accurately, efficiently, and quickly as possible. Any analytical method used should have accuracy, speed, reproducibility, and specificity. No single analytical method is ideally suited for all drugs; each method has its own strengths and weaknesses, and there are a number of factors that determine the validity and reliability of results.

SELECTION OF AN ANALYTICAL METHOD

One general consideration in analytical method selection is the type of information that is needed; quantitative (potency, concentration), semiquantitative (where a “cutoff” level is involved, as in endotoxin levels) or qualitative (yes/no type of testing, including substance identification, sterility). Another consideration involves the physical and chemical characteristics of the analyte, including its solubility, partition coefficient, dissociation constant (pKa), volatility, binding, and the quantity present.

One must consider the degree of quantitative measurement in the validation process, for example accuracy, repeatability/reproducibility, and precision are required; generally, the greater the level that is required, the more sophisticated and expensive the analytical methods that must be used. This is also governed by the types of instrumentation that are on hand or available and the standards available for comparison.

PHYSICAL PHARMACY CAPSULE 4.14 CONT.

FACTORS INVOLVED IN METHODS SELECTION

The ultimate analytical method selected depends upon a number of factors, including sample requirements, sample handling/preparation/purification requirements, type of data needed, and levels of specificity and accuracy required.

SAMPLING REQUIREMENTS

In any analytical method, there may be certain sample requirements that impact one's choice, such as the number of samples needed, the difficulty in obtaining a representative sample, the physical state of the sample (solid, liquid or gas), the type of container required for collection and storage of the sample (some analytes may sorb to the walls or cap liner of the sample containers), and leaching of the container material into the sample, if a liquid, may occur. All these may cause problems in analysis. In the event of sorption, siliconization of the sample vials may sometimes help.

The storage requirements for the sample after collection must be specified (type of container, material used, UV protection, latex contamination, etc.). The effects of air, such as oxidation of the sample ingredients, the presence of carbon dioxide and the formation of insoluble carbonates, pH changes, free versus bound drug, etc., must be considered. The sample must be stored at the proper temperature (refrigerated, frozen or controlled room) prior to shipment and during shipment. Procedures to follow if the sample is accidentally frozen or experiences a freeze-thaw cycle should be detailed.

In considering the chemical and physical stability of the sample, the effects of water must also be considered. If the sample must be maintained in a dry environment, including a desiccant, this should be detailed. The stability of the sample during storage, extraction, and preparation must be determined. The potential for enzymatic breakdown, or other adverse effects of pH, temperature, solvents, bacterial growth, etc., must be addressed. If volatile solvents are required, special handling must be implemented to prevent evaporation because if some of the solvent is allowed to evaporate, the resulting concentration may be falsely elevated.

The sample matrix effects must be determined. Any effects caused by sample viscosity (pipetting, aspiration), ionic strength (immunoassays, dialysis), buffers (ionized/unionized ratio can alter the extraction efficiency of an analyte prior to analysis), and vapor pressure, where drug can be lost must be considered. If any sample pretreatment is required prior to shipment or working in-house, consider any inaccuracies that may occur from pipetting, which is one of the most common sources of analytical errors when working with small volumes.

There must be a consideration of any physical methods of separation and purification that might be used. Most analytical methods require some degree of sample pretreatment to prepare it for analysis. These may include crystallization from solution, distillation, sublimation, solvent extraction, solid-phase extraction, chromatography, centrifugation; the proper choice of separation and purification depends upon the physical and chemical properties of the sample, including its solubility, volatility, binding, quantity present, etc. The effect of any substances in the formulation that may interfere or alter the results must be known beforehand.

DATA INTERPRETATION REQUIREMENTS

The collection of raw data from the analytical process must be done appropriately. One must ensure that appropriate and valid descriptive statistics are used to analyze the data, and that the operating parameters of the analytical instruments are well established. Reference values, if available, should be provided with the analytical results. A description of the analytical controls used by the laboratory is important for documentation, as well as the source of reference standards used to establish standard curves.

PHYSICAL PHARMACY CAPSULE 4.14 CONT.

ANALYTICAL METHODS

In pharmaceutical analysis, analytical methods can be generally divided into physical testing methods, methods that interact with electromagnetic radiation, conductometric techniques, immunoassay methods, separation techniques and others.

Nonspecific methods generally include melting, freezing and boiling points, density, refractive index, polarimetry, ultraviolet/visible spectroscopy, and pH. Methods that are somewhat more specific include infrared spectroscopic, mass spectroscopy, ion selective electrodes, immunoassay methods, and chromatographic methods (high performance liquid chromatography [HPLC] and gas chromatography [GC]), provided proper standards are used.

Methods that can be routinely used for testing incoming bulk materials, whether active or excipients, include melting, freezing and boiling points, density, refractive index, UV/Visible spectroscopy, infrared spectroscopy, polarimetry, pH, and the separation methods. Final products may generally require a method such as HPLC or GC. A classification of analytical methods follows along with suggested analytical methods that can be used for different dosage forms.

CLASSIFICATION OF ANALYTICAL AND MICROBIOLOGICAL METHODS

Physical testing procedures

- Melting point
- Freezing point
- Boiling point
- Density
- Refractive index
- Optical rotation (Polarimetry)
- Thermal analysis
- Color change
- Precipitate formation
- Viscosity change

Interaction of electromagnetic radiation

- Ultraviolet/Visible spectroscopy
- Infrared spectroscopy
- Fluorescence/Phosphorescence spectroscopy
- Raman spectroscopy
- X-ray spectroscopy
- Flame emission and Atomic absorption spectroscopy
- Polarimetry
- Refractometry
- Interferometry

Conductance methods

- pH
- Ion selective electrodes
- Polarography

Immunoassay

- Radioimmunosassay
- Enzyme multiplied immunoassay technique
- Enzyme linked immunosorbent assay
- Fluorescent immunoassay

PHYSICAL PHARMACY CAPSULE 4.14 CONT.

Separation techniques

HPLC

GC

Thin layer chromatography

Paper chromatography

Column chromatography

Gravimetric

Balance

Others

Osmolality

Microbiological methods

Sterility testing

Endotoxin testing

Preservative effectiveness testing

Suggested analytical methods for various dosage forms, depending upon the active drug:

DOSAGE Form	ANALYTICAL METHOD												
	Wt	Vol	pH	Osm	RI	Sp Gr	MP	UV/vis	HPLC	GC	IR	Steril	Endo
Bulk substances	—	—	*	—	*	—	*	*	*	*	*	—	—
Powders	*	—	—	—	—	—	—	—	*	*	—	—	—
Capsules	*	—	—	—	—	—	—	—	*	*	—	—	—
Tablets		*	—	—	—	—	—	—	—	*	*	—	—
Lozenges	*	—	—	—	—	—	—	—	*	*	—	—	—
Suppositories	*	—	—	—	—	*	*	—	*	*	—	—	—
Sticks	*	—	—	—	—	*	*	—	*	*	—	—	—
Solutions	*	*	*	*	*	*	—	*	*	*	—	—	—
Suspensions	*	*	*	—	—	*	—	—	*	*	—	—	—
Emulsions	*	*	*	—	—	*	—	—	*	*	—	—	—
Semisolids	*	—	—	—	—	*	*	—	*	*	—	—	—
Gels	*	*	*	—	*	*	—	—	*	*	—	—	—
Ophthalmics, Otics & Nasals	*	*	*	*	*	*	—	*	*	*	—	*(Oph. only)	—
Inhalations	*	*	*	*	*	—	—	*	*	*	—	*	—
Injections	*	*	*	*	*	*	—	*	*	*	—	*	*

CONSTRUCTION OF A STANDARD CURVE

A standard curve is constructed by analyzing samples (standards) of known composition, generally in increasing concentrations. As each standard is analyzed, an instrumental response (Absorbance, Peak Height, Peak Area, Other Numerical Value) will be obtained. The standard concentrations are plotted as the “x” axis on a graph and the instrumental responses are plotted on the “y” axis. As an example,

The following table represents the results from an HPLC analytical method of methotrexate.

Concentration ($\mu\text{g/mL}$)	0	10	20	30
Response (Peak Height in units)	0	2600	5190	7780

PHYSICAL PHARMACY CAPSULE 4.14 CONT.

When plotted on a graph, one obtains the following:

The next step involves analyzing the unknown sample to obtain a response from the instrument. For example, if the unknown sample provided an instrumental response of 3895, checking that value on the y-axis and moving toward the right on the graph until it intersects the plotted line and dropping down to the x-axis, we can read a value of 15 $\mu\text{g}/\text{mL}$ of the methotrexate. As an option,


the equation of the line can be calculated and the concentration determined by substituting the values of “y” and “b” with the slope of the line to obtain the drug concentration, as follows:

$$m = \Delta y / \Delta x = (7780 - 0) / (30 - 0) = 7780 / 30 = 259.3$$

$$y = mx + b$$

$$3895 = 259.3x + 0$$

$$x = 15.02 \mu\text{g}/\text{mL}$$



ability (lotions), consistency, particle-size distribution, strength, and weight loss.

Ophthalmic and nasal and oral inhalation preparations: Appearance, color, consistency, pH, clarity (solutions), particle size and resuspendability (suspensions, ointments), strength, and sterility.

Small-volume parenterals: Appearance, color, particulate matter, dispersibility (suspensions), pH, sterility, pyrogenicity, and closure integrity.

Large-volume parenterals: Appearance, color, clarity, particulate matter, pH, volume and extractables (when plastic containers are used), sterility, pyrogenicity, and closure integrity.

Suppositories: Softening range, appearance, and melting.

Emulsions: Appearance (such as phase separation), color, odor, pH, and viscosity.

Controlled-release membrane drug delivery systems: Seal strength of the drug reservoir, decomposition products, membrane integrity, drug strength, and drug release rate.

Under usual circumstances, most manufactured products must have a shelf life of 2 or more years to ensure stability at the time of consumption. Commercial products must bear an appropriate expiration date that sets out the time during which the product may be expected to maintain its potency and remain stable under the designated storage conditions. The expiration date limits the time during which the product may be dispensed by the pharmacist or used by the patient.

Prescriptions requiring extemporaneous compounding by the pharmacist do not require the extended shelf life that commercially manufactured and distributed products do because they are intended to be used immediately on receipt by the patient and used only during the immediate course of the prescribed treatment. However, these compounded prescriptions must remain stable and efficacious during the course of use, and the compounding pharmacist must employ formulative components and techniques that will result in a stable product (7).

In years past, pharmacists were confronted primarily with innocuous topical prescriptions that required extemporaneous formulation. However, in recent years there has been a need to compound other drug delivery systems, for example, progesterone vaginal suppositories and oral suspensions, from tablets or capsules. When presented with a prescription that requires extemporaneous compounding, the pharmacist is confronted with a difficult situation, because the potency and stability of these prescriptions is a serious matter. Occasionally, the results of compatibility and stability studies on such prescriptions are published in scientific and professional journals. These are very useful; however, for some prescriptions stability and compatibility information is not readily available. In these instances, it behooves the pharmacist to contact the manufacturer of the active

ingredient or ingredients to solicit stability information. Also, a compilation of published stability information is included in *Trissel's Stability of Compounded Formulations* (8). The published stability data are applicable only to products that are prepared identically to the products that are reported.

USP guidelines on stability of extemporaneous compounded formulations state that in the absence of stability information applicable to a specific drug and preparation, the following guidelines can be used: nonaqueous liquids and solid formulations in which the manufactured drug is the source of the active ingredient, not later than 25% of the time remaining until the product's expiration date or 6 months, whichever is earlier; nonaqueous liquids and solid formulations in which a USP or National Formulary (NF) substance is the source of active ingredient, a beyond-use date of 6 months; for water-containing formulations prepared from ingredients in solid form, a beyond-use date not later than 14 days in storage at cold temperatures; for all other formulations, a beyond-use date of the intended duration of therapy or 30 days, whichever is earlier (7). Thus, if an oral aqueous liquid preparation is made from a tablet or capsule formulation, the pharmacist should make up only at most 14 days' supply, and it must be stored in a refrigerator. Furthermore, the pharmacist must dispense the medication in a container conducive to stability and use and must advise the patient of the proper method of use and conditions of storage of the medication.

Finally, when compounding on the basis of extrapolated or less than concrete information, the pharmacist is well advised to keep the formulation simple and not to shortcut but use the necessary pharmaceutical adjuvants to prepare the prescription.

PHARMACEUTICAL INGREDIENTS AND EXCIPIENTS

DEFINITIONS AND TYPES

To produce a drug substance in a final dosage form requires pharmaceutical ingredients. For example, in the preparation of solutions, one or more *solvents* are used to dissolve the drug substance, *flavors* and *sweeteners* are used to make the product more palatable, *colorants* are added to enhance appeal, *preservatives* may be added

to prevent microbial growth, and *stabilizers*, such as antioxidants and chelating agents, may be used to prevent decomposition, as previously discussed. In the preparation of tablets, *diluents* or *fillers* are commonly added to increase the bulk of the formulation, *binders* to cause adhesion of the powdered drug and pharmaceutical substances, *antiadherents* or *lubricants* to assist smooth tablet formation, *disintegrating agents* to promote tablet breakup after administration, and coatings to improve stability, control disintegration, or enhance appearance. Ointments, creams, and suppositories acquire their characteristic features from their pharmaceutical bases. Thus, for each dosage form, the pharmaceutical ingredients establish the primary features of the product and contribute to the physical form, texture, stability, taste, and overall appearance.

Table 4.3 presents the principal categories of pharmaceutical ingredients, listing some of the official and commercial agents in use. Additional discussion of many ingredients may be found in the chapters where they are most relevant; for example, pharmaceutical materials used in tablet and capsule formulations are discussed in Chapters 7 and 8 and those used in modified-release solid oral dosage forms and drug delivery systems in Chapter 9.

HANDBOOK OF PHARMACEUTICAL EXCIPIENTS AND FOOD AND CHEMICALS CODEX

The *Handbook of Pharmaceutical Excipients* (9) presents monographs on more than 250 excipients used in dosage form preparation. Each monograph includes such information as nonproprietary, chemical, and commercial names; empirical and chemical formulas and molecular weight; pharmaceutical specifications and chemical and physical properties; incompatibilities and interactions with other excipients and drug substances; regulatory status; and applications in pharmaceutical formulation or technology. Additional excipients commonly used are listed in the Food Chemicals Codex (FCC), now owned and published by the USP. The Codex contains information on general provisions and requirements applying to specifications, tests and assays of the FCC, monograph specifications, flavor chemicals, infrared spectra, and general tests and assays.

TABLE 4.3 EXAMPLES OF PHARMACEUTICAL INGREDIENTS

INGREDIENT TYPE	DEFINITION	EXAMPLES
<i>Acidifying agent</i>	Used in liquid preparations to provide acidic medium for product stability	Citric acid Acetic acid Fumaric acid Hydrochloric acid Nitric acid
<i>Alkalinizing agent</i>	Used in liquid preparations to provide alkaline medium for product stability	Ammonia solution Ammonium carbonate Diethanolamine Monoethanolamine Potassium hydroxide Sodium bicarbonate Sodium borate Sodium carbonate Sodium hydroxide Trolamine
<i>Adsorbent</i>	An agent capable of holding other molecules onto its surface by physical or chemical (chemisorption) means	Powdered cellulose Activated charcoal
<i>Aerosol propellant</i>	Agent responsible for developing the pressure within an aerosol container and expelling the product when the valve is opened	Carbon dioxide Dichlorodifluoromethane Dichlorotetrafluoroethane Trichloromonofluoromethane
<i>Air displacement</i>	Agent employed to displace air in a hermetically sealed container to enhance product stability	Nitrogen Carbon dioxide
<i>Antifungal preservative</i>	Used in liquid and semisolid preparations to prevent growth of fungi. Effectiveness of parabens is usually enhanced by use in combination	Butylparaben Ethylparaben Methylparaben Benzoic acid Propylparaben Sodium benzoate Sodium propionate
<i>Antimicrobial preservative</i>	Used in liquid and semisolid preparations to prevent growth of microorganisms	Benzalkonium chloride
<i>Antioxidant</i>	Used to prevent deterioration of preparations by oxidation	Ascorbic acid Ascorbyl palmitate Butylated hydroxyanisole Butylated hydroxytoluene Hypophosphorous acid Monothioglycerol Propyl gallate Sodium ascorbate Sodium bisulfite Sodium formaldehyde Sulfoxylate Sodium metabisulfite
<i>Buffering agent</i>	Used to resist change in pH upon dilution or addition of acid or alkali	Potassium metaphosphate Potassium phosphate, monobasic Sodium acetate Sodium citrate, anhydrous and dihydrate
<i>Chelating agent</i>	Substance that forms stable water-soluble complexes (chelates) with metals; used in some liquid pharmaceuticals as stabilizers to complex heavy metals that might promote instability. In such use, they are also called <i>sequestering agents</i>	Edetic acid Edetate disodium

(continued)

TABLE 4.3 EXAMPLES OF PHARMACEUTICAL INGREDIENTS (Continued)

INGREDIENT TYPE	DEFINITION	EXAMPLES
<i>Colorant</i>	Used to impart color to liquid and solid (e.g., tablets and capsules) preparations	FD&C Red No. 3 FD&C Red No. 20 FD&C Yellow No. 6 FD&C Blue No. 2 D&C Green No. 5 D&C Orange No. 5 D&C Red No. 8 Caramel Ferric oxide, red
<i>Clarifying agent</i>	Used as a filtering aid for its adsorbent qualities	Bentonite
<i>Emulsifying agent</i>	Used to promote and maintain dispersion of finely subdivided particles of liquid in a vehicle in which it is immiscible. End product may be a liquid emulsion or semisolid emulsion (e.g., a cream)	Acacia Cetomacrogol Cetyl alcohol Glyceryl monostearate Sorbitan monooleate Polyoxyethylene 50 stearate
<i>Encapsulating agent</i>	Used to form thin shells to enclose a drug for ease of administration	Gelatin
<i>Flavorant</i>	Used to impart a pleasant flavor and often odor to a preparation. In addition to the natural flavorants listed, many synthetic ones are used	Anise oil Cinnamon oil Cocoa Menthol Orange oil Peppermint oil Vanillin
<i>Humectant</i>	Used to prevent drying of preparations, particularly ointments and creams	Glycerin Propylene glycol Sorbitol
<i>Levigating agent</i>	Liquid used as an intervening agent to reduce the particle size of a powder by grinding, usually in a mortar	Mineral oil Glycerin Propylene glycol
<i>Ointment base</i>	Semisolid vehicle for medicated ointments	Lanolin Hydrophilic ointment Polyethylene glycol ointment Petrolatum Hydrophilic petrolatum White ointment Yellow ointment Rose water ointment
<i>Plasticizer</i>	Component of film-coating solutions to make film more pliable, enhance spread of coat over tablets, beads, and granules	Diethyl phthalate Glycerin
<i>Solvent</i>	Used to dissolve another substance in preparation of a solution; may be aqueous or not (e.g., oleaginous). Cosolvents, such as water and alcohol (hydroalcoholic) and water and glycerin, may be used when needed. Sterile solvents are used in certain preparations (e.g., injections)	Alcohol Corn oil Cottonseed oil Glycerin Isopropyl alcohol Mineral oil Oleic acid Peanut oil Purified water Water for injection Sterile water for injection Sterile water for irrigation

(continued)

TABLE 4.3 EXAMPLES OF PHARMACEUTICAL INGREDIENTS (Continued)

INGREDIENT TYPE	DEFINITION	EXAMPLES
<i>Stiffening agent</i>	Used to increase thickness or hardness of a preparation, usually an ointment	Cetyl alcohol Cetyl esters wax Microcrystalline wax Paraffin Stearyl alcohol White wax Yellow wax
<i>Suppository base</i>	Vehicle for suppositories	Cocoa butter Polyethylene glycols (mixtures) PEG 3350
<i>Surfactant (surface active agent)</i>	Substances that absorb to surfaces or interfaces to reduce surface or interfacial tension. May be used as wetting agents, detergents, or emulsifying agents	Benzalkonium chloride Nonoxynol 10 Octoxynol 9 Polysorbate 80 Sodium lauryl sulfate Sorbitan monopalmitate
<i>Suspending agent</i>	Viscosity-increasing agent used to reduce sedimentation rate of particles in a vehicle in which they are not soluble; suspension may be formulated for oral, parenteral, ophthalmic, topical, or other route	Agar Bentonite Carbomer (e.g., Carbopol) Carboxymethylcellulose sodium Hydroxyethyl cellulose Hydroxypropyl cellulose Hydroxypropyl methylcellulose Kaolin Methylcellulose Tragacanth Veegum
<i>Sweetening agent</i>	Used to impart sweetness to a preparation	Aspartame Dextrose Glycerin Mannitol Saccharin sodium Sorbitol Sucrose
<i>Tablet antiadherents</i>	Prevent tablet ingredients from sticking to punches and dies during production	Magnesium stearate
<i>Tablet binders</i>	Substances used to cause adhesion of powder particles in tablet granulations	Acacia Alginic acid Carboxymethylcellulose sodium Compressible sugar (e.g., Nu-Tab) Ethylcellulose Gelatin Liquid glucose Methylcellulose Povidone Pregelatinized starch
<i>Tablet and capsule diluent</i>	Inert filler to create desired bulk, flow properties, and compression characteristics of tablets and capsules	Dibasic calcium phosphate Kaolin Lactose Mannitol Microcrystalline cellulose Powdered cellulose Precipitated calcium carbonate Sorbitol Starch

(continued)

TABLE 4.3 EXAMPLES OF PHARMACEUTICAL INGREDIENTS (Continued)

INGREDIENT TYPE	DEFINITION	EXAMPLES
<i>Tablet coating agent</i>	Used to coat a tablet to protect against decomposition by atmospheric oxygen or humidity, to provide a desired release pattern, to mask taste or odor, or for aesthetic purposes. Coating may be sugar, film, or thick covering around a tablet. Sugar-coated tablets generally start to break up in the stomach. Film forms a thin cover around a formed tablet or bead. Unless it is enteric, film dissolves in the stomach. Enteric coating passes through the stomach to break up in the intestines. Some water-insoluble coatings (e.g., ethylcellulose) are used to slow the release of drug in the gastrointestinal tract	
<i>Sugar coating</i>		Liquid glucose Sucrose
<i>Film coating</i>		Hydroxyethyl cellulose Hydroxypropyl cellulose Hydroxypropyl methylcellulose Methylcellulose (e.g., Methocel) Ethylcellulose (e.g., Ethocel)
<i>Enteric coating</i>		Cellulose acetate phthalate Shellac (35% in alcohol, pharmaceutical glaze)
<i>Tablet direct compression excipient</i>	Used in direct compression tablet formulations	Dibasic calcium phosphate (e.g., Datab)
<i>Tablet disintegrant</i>	Used in solid forms to promote disruption of the mass into smaller particles more readily dispersed or dissolved	Alginate acid Polacrillin potassium (e.g., Amberlite) Sodium alginate Sodium starch glycolate Starch
<i>Tablet glidant</i>	Used in tablet and capsule formulations to improve flow properties of the powder mixture	Colloidal silica Cornstarch Talc
<i>Tablet lubricant</i>	Used in tablet formulations to reduce friction during tablet compression	Calcium stearate Magnesium stearate Mineral oil Stearic acid Zinc stearate
<i>Tablet or capsule opaquant</i>	Used to render a coating opaque. May be used alone or with a colorant	Titanium dioxide
<i>Tablet polishing agent</i>	Used to impart an attractive sheen to coated tablets	Carnauba wax White wax
<i>Tonicity agent</i>	Used to render solution similar in osmotic-dextrose characteristics to physiologic fluids, e.g., in ophthalmic, parenteral, and irrigation fluids	Sodium chloride
<i>Vehicle</i>	Carrying agent used in formulating a variety of liquids for oral and parenteral administration Generally, oral liquids are aqueous (e.g., syrups) or hydroalcoholic (e.g., elixirs). Solutions for intravenous use are aqueous, whereas intramuscular injections may be aqueous or oleaginous	

(continued)

TABLE 4.3 EXAMPLES OF PHARMACEUTICAL INGREDIENTS (Continued)

INGREDIENT TYPE	DEFINITION	EXAMPLES
<i>Flavored, sweetened</i>		Acacia syrup Aromatic syrup Aromatic elixir Cherry syrup Cocoa syrup Orange syrup Syrup
<i>Oleaginous</i>		Corn oil Mineral oil Peanut oil Sesame oil
<i>Sterile</i>		Bacteriostatic sodium chloride injection
<i>Viscosity-increasing agent</i>	Used to render preparations more resistant to flow. Used in suspensions to deter sedimentation, in ophthalmic solutions to enhance contact time (e.g., methylcellulose), to thicken topical creams, etc.	Alginic acid Bentonite Carbomer Carboxymethylcellulose Sodium Methylcellulose Povidone Sodium alginate Tragacanth

HARMONIZATION OF STANDARDS

There is great interest in the international harmonization of standards applicable to pharmaceutical excipients. This is because the pharmaceutical industry is multinational, with major companies having facilities in more than a single country, with products sold in markets worldwide, and with regulatory approval for these products required in each country. Standards for each drug substance and excipient used in pharmaceuticals are contained in pharmacopeias—or for new agents, in an application for regulatory approval by the relevant governing authority. The four pharmacopeias with the largest international use are the *United States Pharmacopeia–National Formulary* (USP–NF), *British Pharmacopeia*, *European Pharmacopeia*, and *Japanese Pharmacopeia*. Uniform standards for excipients in these and other pharmacopeias would facilitate production efficiency, enable the marketing of a single formulation of a product internationally, and enhance regulatory approval of pharmaceutical products worldwide. The goal of harmonization is an ongoing effort by corporate representatives and international regulatory authorities.

A few of the more common and widely used pharmaceutical excipients, including sweeteners,

flavors, colors, and preservatives, are discussed here.

APPEARANCE AND PALATABILITY

Although most drug substances in use today are unpalatable and unattractive in their natural state, their preparations present them to the patient as colorful, flavorful formulations attractive to the sight, smell, and taste. These qualities, which are the rule rather than the exception, have virtually eliminated the natural reluctance of many patients to take medications because of disagreeable odor or taste. In fact, the inherent attractiveness of today's pharmaceuticals has caused them to acquire the dubious distinction of being a source of accidental poisonings in the home, particularly among children who are lured by their organoleptic appeal.

There is some psychologic basis to drug therapy, and the odor, taste, and color of a pharmaceutical preparation can play a part. An appropriate drug has its most beneficial effect when it is accepted and taken properly by the patient. The proper combination of flavor, fragrance, and color in a pharmaceutical product contributes to its acceptance.




FIGURE 4.4 Electronic tongue to assist in formulation development. (Courtesy of Alpha MOS.)

An “electronic tongue” is used to aid in providing a global “taste fingerprint” during formulation development. It provides information on bitterness levels and the stability of flavors in terms of taste (Figure 4.4).

Flavoring Pharmaceuticals

The flavoring of pharmaceuticals applies primarily to liquids intended for oral administration. The 10,000 taste buds on the tongue, roof of the mouth, cheeks, and throat have 60 to 100 receptor cells each (10). These receptor cells interact with molecules dissolved in the saliva and produce a positive or negative taste sensation. Medication in liquid form comes into immediate and direct contact with these taste buds. The addition of flavoring agents to liquid medication can mask the disagreeable taste. Drugs placed in capsules or prepared as coated tablets may be easily swallowed with no contact between the drug and the taste buds. Tablets containing drugs that are not especially distasteful may remain uncoated and unflavored. Swallowing them with water usually is sufficient to avoid undesirable taste sensations. However, chewable tablets, such as certain antacid and vitamin products, usually are sweetened and flavored to improve acceptance.

The flavor sensation of a food or pharmaceutical is actually a complex blend of taste and smell, with lesser influences of texture, temperature, and even sight. In flavor-formulating a pharmaceutical product, the pharmacist must give consideration to the color, odor, texture, and taste of the preparation. It would be incongruous, for example, to color a liquid pharmaceutical

red and give it a banana taste and a mint odor. The color of a pharmaceutical must have a psychogenic balance with the taste, and the odor must also enhance that taste. Odor greatly affects the flavor of a preparation or foodstuff. If one’s sense of smell is impaired, as during a head cold, the usual flavor sensation of food is similarly diminished.

The medicinal chemist and the formulation pharmacist are well acquainted with the taste characteristics of certain chemical types of drugs and strive to mask the unwanted taste through the appropriate use of flavoring agents. Although there are no rules for unerringly predicting the taste sensation of a drug based on its chemical constitution, experience permits the presentation of several observations. For instance, although we recognize and assume the salty taste of sodium chloride, the formulation pharmacist knows that not all salts are salty but that their taste is a function of both cation and anion. Whereas salty tastes are evoked by chlorides of sodium, potassium, and ammonium and by sodium bromide, bromides of potassium and ammonium elicit bitter and salty sensations, and potassium iodide and magnesium sulfate (Epsom salt) are predominantly bitter. In general, low-molecular-weight salts are salty, and high-molecular-weight salts are bitter.

With organic compounds, an increase in the number of hydroxyl groups ($-\text{OH}$) seems to increase the sweetness of the compound. Sucrose, which has eight hydroxyl groups, is sweeter than glycerin, another pharmaceutical sweetener, which has but three hydroxyl groups. In general, the organic esters, alcohols, and aldehydes are pleasant to the taste, and since many of them are volatile, they also contribute to the odor and thus the flavor of preparations in which they are used. Many nitrogen-containing compounds, especially the plant alkaloids (e.g., quinine) are extremely bitter, but certain other nitrogen-containing compounds (e.g., aspartame) are extremely sweet. The medicinal chemist recognizes that even the most simple structural change in an organic compound can alter its taste. D-Glucose is sweet, but L-glucose has a slightly salty taste; saccharin is very sweet, but N-methyl-saccharin is tasteless (11).

Thus, prediction of the taste characteristics of a new drug is only speculative. However, it is soon learned and the formulation pharmacist is

then put to the task of increasing the drug's palatability in the environment of other formulative agents. The selection of an appropriate flavoring agent depends on several factors, primarily the taste of the drug substance itself. Certain flavoring materials are more effective than others in masking or disguising the particular bitter, salty, sour, or otherwise undesirable taste of medicinal agents. Although individuals' tastes and flavor preferences differ, cocoa-flavored vehicles are considered effective for masking the taste of bitter drugs. Fruit or citrus flavors are frequently used to combat sour or acid-tasting drugs, and cinnamon, orange, raspberry, and other flavors have been successfully used to make preparations of salty drugs more palatable.

The age of the intended patient should also be considered in the selection of the flavoring agent, because certain age groups seem to prefer certain flavors. Children prefer sweet candy-like preparations with fruity flavors, but adults seem to prefer less sweet preparations with a tart rather than a fruit flavor.

Flavors can consist of oil- or water-soluble liquids and dry powders; most are diluted in carriers. Oil-soluble carriers include soybean and other edible oils; water-soluble carriers include water, ethanol, propylene glycol, glycerin, and emulsifiers. Dry carriers include maltodextrins, corn syrup solids, modified starches, gum arabic, salt, sugars, and whey protein. Flavors can degrade as a result of exposure to light, temperature, head space oxygen, water, enzymes, contaminants, and other product components, so they must be carefully selected and checked for stability.

The different types of flavors include natural, artificial, and spice:

Natural flavor: Essential oil, oleoresin, essence or extractive, protein hydrolysate, distillate, or any product of roasting, heating, or enzymolysis, which contains the flavoring constituents derived from a spice, fruit or fruit juice, vegetable or vegetable juice, edible yeast, herb, bark, bud, root, leaf or similar plant material, meat, seafood, poultry, eggs, dairy products, or fermentation products thereof whose significant function in food is flavoring rather than nutritional. [CFR 101.22(a)(3)] In "all natural" flavors, one doesn't necessarily know the exact chemical composition.

Artificial flavor: Any substance used to impart flavor that is not derived from a spice, fruit or fruit juice, vegetable or vegetable juice, edible yeast, herb, bark, bud, root, leaf or similar plant material, meat, fish, poultry, eggs, dairy products, or fermentation products thereof. [CFR 101.22(a)(1)]

Spice: Any aromatic vegetable substance in whole, broken, or ground form, except substances traditionally regarded as foods, such as onions, garlic, and celery; whose significant function in food is seasoning rather than nutritional; that is true to name; and from which no portion of any volatile oil or other flavoring principle has been removed. [CFR 101.22(a)(2)]

In addition to the types of flavors, you should be aware of commercial flavor designations, including the following (*Note:* ABCD would be the flavor name, e.g., cherry):

Natural ABCD flavor	All components derived from ABCD.
ABCD flavor, natural	At least one component and artificial-derived from ABCD. No definition of natural to artificial ratio.
ABCD flavor, WONF*	All components natural. At least one component derived from ABCD.
Natural flavor, ABCD type	All components natural. No components derived from ABCD.
ABCD flavor, artificial	All components are artificial.
Conceptual flavors	May contain artificial flavors. No reference point. May only have to declare in ingredient declaration.

*WONF, with other natural flavors.

A general guide to using flavors is to start as follows (keep in mind it is usually possible to add more flavor, but once it is added, it is too late to remove it).

Water-soluble flavors	Generally start at 0.2% for artificial and 1%–2% for natural flavors.
Oil-soluble flavors	Generally start at 0.1% in finished product for artificial flavors and 0.2% for natural flavors.
Powdered flavors	Generally start at 0.1% in finished product for artificial flavors and 0.75% for natural flavors.

Sweetening Pharmaceuticals

In addition to sucrose, a number of artificial sweetening agents have been used in foods and pharmaceuticals over the years. Some of these, including aspartame, saccharin, and cyclamate, have faced challenges over their safety by the FDA and restrictions to their use and sale; in fact, in 1969, FDA banned cyclamates from use in the United States.

The introduction of diet soft drinks in the 1950s provided the spark for the widespread use of artificial sweeteners today. Besides dieters, patients with diabetes are regular users of artificial sweeteners. Over the years, each of the artificial sweeteners has undergone long periods of review and debate. Critical to the evaluation of food additives are issues of metabolism and toxicity. For example, almost none of the saccharin a person consumes is metabolized; it is excreted by the kidneys virtually unchanged. Cyclamate, on the other hand, is metabolized, or processed, in the digestive tract, and its by-products are excreted by the kidneys. Aspartame breaks down in the body into three basic components: the amino acids phenylalanine and aspartic acid, and methanol. These three components, which also occur naturally in various foods, are in turn metabolized through regular pathways in the body. Because of its metabolism to phenylalanine, the use of aspartame by persons with phenylketonuria (PKU) is discouraged, and diet foods and drinks must bear an appropriate label warning indicating that the particular foodstuff not be consumed by such individuals. They cannot metabolize phenylalanine adequately, so they undergo an increase in the serum levels of the amino acid (hyperphenylalaninemia). This can result in mental retardation and can affect the fetus of a pregnant woman who has PKU.

Passage in 1958 of the Food Additives Amendment to the Food, Drug, and Cosmetic Act produced a major change in how the federal government regulates food additives. For one thing, no new food additive may be used if animal feeding studies or other appropriate tests showed that it caused cancer. This is the famous Delaney Clause. The *amount* of the substance one would have to consume to induce cancer is not significant under the Delaney Clause.

Another critical feature of the 1958 amendment was that it did not apply to additives that were generally recognized by experts as safe for their intended uses. Saccharin, cyclamate, and a long list of other substances were being used in foods before the amendment's passage and were "generally recognized as safe"—or what is known today as GRAS. Aspartame, on the other hand, was the first artificial sweetener to fall under the 1958 amendment's requirement for premarketing proof of safety, because the first petition to FDA for its approval was filed in 1973. In 1968, the Committee on Food Protection of the National

Academy of Sciences issued an interim report on the safety of nonnutritive sweeteners, including saccharin. In the early 1970s, FDA began a major review of hundreds of food additives on the GRAS list to determine whether current studies justified their safe status. In 1972, with new studies under way, FDA decided to take saccharin off the GRAS and establish interim limits that would permit its continued use until additional studies were completed. (Previous studies indicated that male and female rats fed doses of saccharin developed a significant incidence of bladder tumors.) In November 1977, Congress passed the Saccharin Study and Labeling Act, which permitted saccharin's continued availability while mandating that warning labels be used to advise consumers that saccharin caused cancer in animals. The law also directed FDA to arrange further studies of carcinogens and toxic substances in foods.

Cyclamate was introduced into beverages and foods in the 1950s and dominated the artificial sweetener market in the 1960s. After much controversy regarding its safety, the FDA issued a final ruling in 1980 stating that safety has not been demonstrated. Since that date, scientific studies have continued the search for conclusive support or rejection of the FDA decision. At question is cyclamate's possible carcinogenicity and its possible causation of genetic damage and testicular atrophy. See the indicated references for a review of the recent history of sweeteners, including saccharin, cyclamate, fructose, polyalcohols, sucrose, and aspartame (12–15).

Acesulfame potassium, a nonnutritive sweetener discovered in 1967, was approved in 1992 by the FDA. It previously was used in a number of other countries. Structurally similar to saccharin, it is 130 times as sweet as sucrose and is excreted unchanged in the urine. Acesulfame is more stable than aspartame at elevated temperatures and FDA initially approved it for use in candy, chewing gum, confectionery, and instant coffee and tea.

A relatively new sweetening agent in U.S. commerce is Stevia powder, the extract from the leaves of the plant *Stevia rebaudiana bertonii*. It is natural, nontoxic, safe, and about 30 times as sweet as cane sugar, or sucrose. It can be used in both hot and cold preparations. Table 4.4 compares three of the most commonly used sweeteners in the food and drug industry.

Most large pharmaceutical manufacturers have special laboratories for taste-testing

TABLE 4.4 COMPARISON OF SWEETENERS

	SUCROSE	SACCHARIN	ASPARTAME
Source	Sugar cane; sugar beet	Chemical synthesis; phthalic anhydride, a petroleum product	Chemical synthesis; methyl ester dipeptide of phenylalanine and aspartic acid
Relative	1	300	180–200 sweetness
Bitterness	None	Moderate to strong	None
Aftertaste	None	Moderate to strong; sometimes metallic or bitter	None
Calories	4/g	0	4/g
Acid stability	Good	Excellent	Fair
Heat stability	Good	Excellent	Poor

proposed formulations of their products. Panels of employees or interested community participants participate in evaluating the various formulations, and their assessments become the basis for the firm's flavoring decisions.

The flavoring agent in liquid pharmaceutical products is added to the solvent or vehicle component of the formulation in which it is most soluble or miscible. That is, water-soluble flavorants are added to the aqueous component of a formulation and poorly water-soluble flavorants are added to the alcoholic or other non-aqueous solvent component of the formulation. In a hydroalcoholic or other multisolvent system, care must be exercised to maintain the flavorant in solution. This is accomplished by maintaining a sufficient level of the flavorant's solvent.

Coloring Pharmaceuticals

Coloring agents are used in pharmaceutical preparations for esthetics. A distinction should be made between agents that have inherent color and those that are employed as colorants. Certain agents—sulfur (yellow), riboflavin (yellow), cupric sulfate (blue), ferrous sulfate (bluish green), cyanocobalamin (red), and red mercuric iodide (vivid red)—have inherent color and are not thought of as pharmaceutical colorants in the usual sense of the term.

Although most pharmaceutical colorants in use today are synthetic, a few are obtained from natural mineral and plant sources. For example, red ferric oxide is mixed in small proportions with zinc oxide powder to give calamine its characteristic pink color, which is intended to match the skin tone upon application.

The synthetic coloring agents used in pharmaceutical products were first prepared in the middle of the 19th century from principles of coal tar. Coal tar (*pix carbonis*), a thick, black viscid liquid, is a by-product of the destructive distillation of coal. Its composition is extremely complex, and many of its constituents may be separated by fractional distillation. Among its products are anthracene, benzene, naphtha, creosote, phenol, and pitch. About 90% of the dyes used in the products FDA regulates are synthesized from a single colorless derivative of benzene called aniline. These aniline dyes are also known as synthetic organic dyes or as coal tar dyes, since aniline was originally obtained from bituminous coal. Aniline dyes today come mainly from petroleum.

Many coal tar dyes were originally used indiscriminately in foods and beverages to enhance their appeal without regard to their toxic potential. It was only after careful scrutiny that some dyes were found to be hazardous to health because of either their own chemical nature or the impurities they carried. As more dyestuffs became available, some expert guidance and regulation were needed to ensure the safety of the public. After passage of the Food and Drug Act in 1906, the U.S. Department of Agriculture established regulations by which a few colorants were *permitted* or *certified* for use in certain products. Today, the FDA regulates the use of color additives in foods, drugs, and cosmetics through the provisions of the Federal Food, Drug, and Cosmetic Act of 1938, as amended in 1960 with the Color Additive Amendments. Lists of color additives *exempt* from certification and those *subject* to certification are codified into law and regulated by the FDA (16). Certified color

additives are classified according to their approved use: (a) FD&C color additives, which may be used in foods, drugs, and cosmetics; (b) D&C color additives, some of which are approved for use in drugs, some in cosmetics, and some in medical devices; and (c) external D&C color additives, the use of which is restricted to external parts of the body, not including the lips or any other body surface covered by mucous membrane. Each certification category has a variety of basic colors and shades for coloring pharmaceuticals. One may select from a variety of FD&C, D&C, and external D&C reds, yellows, oranges, greens, blues, and violets. By selective combinations of the colorants one can create distinctive colors (Table 4.5).

TABLE 4.5 EXAMPLES OF COLOR FORMULATIONS

SHADE OR COLOR	FD&C DYE	% OF BLEND
Orange	Yellow No. 6 or	100
	Yellow No. 5	95
	Red No. 40	5
Cherry	Red No. 40 or	100
	Red No. 40	99
	Blue No. 1	1
Strawberry	Red No. 40 or	100
	Red No. 40	95
	Red No. 3	5
Lemon	Yellow No. 5	100
Lime	Yellow No. 5	95
	Blue No. 1	5
Grape	Red No. 40	80
	Blue No. 1	20
Raspberry	Red No. 3	75
	Yellow No. 6	20
	Blue No. 1	5
Butterscotch	Yellow No. 5	74
	Red No. 40	24
	Blue No. 1	2
Chocolate	Red No. 40	52
	Yellow No. 5	40
	Blue No. 1	8
Caramel	Yellow No. 5	64
	Red No. 3	21
	Yellow No. 6	9
	Blue No. 1	6
Cinnamon	Yellow No. 5	60
	Red No. 40	35
	Blue No. 1	5

From literature of Warner-Jenkinson Co., St. Louis, Mo.

As a part of the National Toxicology Program of the U.S. Department of Health and Human Services, various substances, including color additives, are studied for toxicity and carcinogenesis. For color additives, the study protocols usually call for a 2-year study in which groups of male and female mice and rats are fed diets containing various quantities of the colorant. The killed and surviving animals are examined for evidence of long-term toxicity and carcinogenesis. Five categories of evidence of carcinogenic activity are used in reporting observations: (a) “clear evidence” of carcinogenic activity; (b) “some evidence”; (c) “equivocal evidence,” indicating uncertainty; (d) “no evidence,” indicating no observable effect; and (e) “inadequate study,” for studies that cannot be evaluated because of major flaws.

The certification status of the colorants is continually reviewed, and changes are made in the list of certified colors in accordance with toxicology findings. These changes may be (a) the withdrawal of certification, (b) the transfer of a colorant from one certification category to another, or (c) the addition of new colors to the list. Before gaining certification, a color additive must be demonstrated to be safe. In the case of pharmaceutical preparations, color additives, as with all additives, must not interfere with therapeutic efficacy, nor may they interfere with the prescribed assay procedure for the preparation.

In the 1970s, concern and scientific questioning of the safety of some color additives heightened. A color that drew particular attention was FD&C Red No. 2, because of its extensive use in foods, drugs, and cosmetics. Researchers in Russia reported that this color, also known as amaranth, caused cancer in rats. Although the FDA was never able to determine the purity of the amaranth tested in Russia, these reports led to FDA investigations and a series of tests that eventually resulted in withdrawal of FD&C Red No. 2 from the FDA certified list in 1976 because its sponsors were unable to prove safety. That year, FDA also terminated approval for use of FD&C Red No. 4 in maraschino cherries and ingested drugs because of unresolved safety questions. FD&C Red No. 4 is now permitted only in externally applied drugs and cosmetics.

FD&C Yellow No. 5 (also known as tartrazine) causes allergic-type reactions in many people. People who are allergic to aspirin are also

likely to be allergic to this dye. As a result, the FDA requires listing of this dye by name on the labels of foods (e.g., butter, cheese, ice cream) and ingested drugs containing it.

A colorant becomes an integral part of a pharmaceutical formulation, and its exact quantitative amount must be reproducible each time the formulation is prepared, or else the preparation would have a different appearance from batch to batch. This requires a high degree of skill, for the amount of colorant generally added to liquid preparations ranges from 0.0005% to 0.001% depending upon the colorant and the depth of color desired. Because of their color potency, dyes generally are added to pharmaceutical preparations in the form of diluted solutions rather than as concentrated dry powders. This permits greater accuracy in measurement and more consistent color production.

In addition to liquid dyes in the coloring of pharmaceuticals, lake pigments may also be used. Whereas a chemical material exhibits coloring power or tinctorial strength when dissolved, pigment is an insoluble material that colors by dispersion. An FD&C lake is a pigment consisting of a substratum of alumina hydrate on which the dye is adsorbed or precipitated. Having aluminum hydroxide as the substrate, the lakes are insoluble in nearly all solvents. FD&C lakes are subject to certification and must be made from certified dyes. Lakes do not have a specified dye content; they range from 10% to 40% pure dye. By their nature, lakes are suitable for coloring products in which the moisture levels are low.

Lakes in pharmaceuticals are commonly used in the form of fine dispersions or suspensions. The pigment particles may range in size from less than 1 μm up to 30 μm . The finer the particle, the less chance for color speckling in the finished product. Blends of various lake pigments may be used to achieve a variety of colors, and various vehicles, such as glycerin, propylene glycol, and sucrose-based syrup, may be employed to disperse the colorants.

Colored empty gelatin capsule shells may be used to hold a powdered drug mixture. Many commercial capsules are prepared with a capsule body of one color and a cap of a different color, resulting in a two-colored capsule. This makes certain commercial products more readily identifiable than solid-colored capsules. For powdered drugs dispensed as such or compressed into tab-

lets, a generally larger proportion of dye is required (about 0.1%) to achieve the desired hue than with liquid preparations.

Both dyes and lakes are used to color sugar-coated tablets, film-coated tablets, direct-compression tablets, pharmaceutical suspensions, and other dosage forms (17). Traditionally, sugar-coated tablets have been colored with syrup solutions containing varying amounts of the water-soluble dyes, starting with very dilute solutions, working up to concentrated color syrup solutions. As many as 30 to 60 coats are common. With the lakes, fewer color coats are used. Appealing tablets have been made with as few as 8 to 12 coats using lakes dispersed in syrup. Water-soluble dyes in aqueous vehicles or lakes dispersed in organic solvents may be effectively sprayed on tablets to produce attractive film coatings. There is continued interest today in chewable tablets, because of the availability of many direct-compression materials such as dextrose, sucrose, mannitol, sorbitol, and spray-dried lactose. The direct-compression colored chewable tablets may be prepared with 1 lb of lake per 1,000 lb of tablet mix. For aqueous suspensions, FD&C water-soluble colors or lakes may be satisfactory. In other suspensions, FD&C lakes are necessary. The lakes, added to either the aqueous or nonaqueous phase, generally at a level of 1 lb of color per 1,000 lb of suspension, require homogenization or mechanical blending to achieve uniform coloring.

For the most part, ointments, suppositories, and ophthalmic and parenteral products assume the color of their ingredients and do not contain color additives. Should a dye lose the certification status it held when a product was first formulated, manufactured, and marketed, the manufacturer must reformulate within a reasonable length of time, using only color additives certified at the new date of manufacture.

In addition to esthetics and the certification status of a dye, a formulation pharmacist must select the dyes to be used in a particular formula on the basis of their physical and chemical properties. Of prime importance is the solubility of a prospective dye in the vehicle to be used for a liquid formulation or in a solvent to be employed during a pharmaceutical process, as when the dye is sprayed on a batch of tablets. In general, most dyes are broadly grouped into those that are water soluble and those that are oil soluble; few if any dyes are both. Usually, a water-soluble

dye is also adequately soluble in commonly used pharmaceutical liquids like glycerin, alcohol, and glycol ethers. Oil-soluble dyes may also be soluble to some extent in these solvents and in liquid petrolatum (mineral oil), fatty acids, fixed oils, and waxes. No great deal of solubility is required, since the concentration of dye in a given preparation is minimal.

Another important consideration when selecting a dye for use in a liquid pharmaceutical is the pH and pH stability of the preparation to be colored. Dyes can change color with a change in pH, and the dye must be selected so that no anticipated pH change will alter the color during the usual shelf life. The dye also must be chemically stable in the presence of the other formula-tive ingredients and must not interfere with the stability of the other agents. To maintain their original colors, FD&C dyes must be protected from oxidizing agents, reducing agents (especially metals, including iron, aluminum, zinc, and tin), strong acids and alkalis, and excessive heating. Dyes must also be reasonably photo-stable; that is, they must not change color when exposed to light of anticipated intensities and wavelengths under the usual conditions of shelf storage. Certain medicinal agents, particularly those prepared in liquid form, must be protected from light to maintain their chemical stability and their therapeutic effectiveness. These preparations are generally kept in dark amber or opaque containers. For solid dosage forms of photolabile drugs, a colored or opaque capsule shell may enhance the drug's stability by shielding out light rays.

PRESERVATIVES

In addition to the stabilization of pharmaceutical preparations against chemical and physical degradation due to changed environmental conditions within a formulation, certain liquid and semisolid preparations must be preserved against microbial contamination.

Sterilization and Preservation

Although some types of pharmaceutical products, for example, ophthalmic and injectable preparations, are sterilized by physical methods (autoclaving for 20 minutes at 15 lb pressure and 121°C, dry heat at 180°C for 1 hour, or bacterial filtration) during manufacture, many of them

also require an antimicrobial preservative to maintain their aseptic condition throughout storage and use. Other types of preparations that are not sterilized during their preparation but are particularly susceptible to microbial growth because of the nature of their ingredients are protected by the addition of an antimicrobial preservative. Preparations that provide excellent growth media for microbes are most aqueous preparations, especially syrups, emulsions, suspensions, and some semisolid preparations, particularly creams. Certain hydroalcoholic and most alcoholic preparations may not require the addition of a chemical preservative when the alcoholic content is sufficient to prevent microbial growth. Generally, 15% V/V alcohol will prevent microbial growth in acid media and 18% V/V in alkaline media. Most alcohol-containing pharmaceuticals, such as elixirs, spirits, and tinctures, are self-sterilizing and do not require additional preservation. The same applies to other individual pharmaceuticals that by virtue of their vehicle or other formula-tive agents may not permit the growth of microorganisms.

Preservative Selection

When experience or shelf storage experiments indicate that a preservative is required in a pharmaceutical preparation, its selection is based on many considerations, including some of the following:

- The preservative prevents the growth of the type of microorganisms considered the most likely contaminants of the preparation.
- The preservative is soluble enough in water to achieve adequate concentrations in the aqueous phase of a system with two or more phases.
- The proportion of preservative remaining undissociated at the pH of the preparation makes it capable of penetrating the microorganism and destroying its integrity.
- The required concentration of the preservative does not affect the safety or comfort of the patient when the pharmaceutical preparation is administered by the usual or intended route; that is, it is nonirritating, nonsensitizing, and nontoxic.
- The preservative has adequate stability and will not be reduced in concentration by chemical decomposition or volatilization during the desired shelf life of the preparation.

- The preservative is completely compatible with all other formulative ingredients and does not interfere with them, nor do they interfere with the effectiveness of the preservative agent.
- The preservative does not adversely affect the preparation's container or closure.

General Preservative Considerations

Microorganisms include molds, yeasts, and bacteria, with bacteria generally favoring a slightly alkaline medium and the others an acid medium. Although few microorganisms can grow below pH 3 or above pH 9, most aqueous pharmaceutical preparations are within the favorable pH range and therefore must be protected against microbial growth. To be effective, a preservative agent must be dissolved in sufficient concentration in the aqueous phase of a preparation. Furthermore, only the undissociated fraction or molecular form of a preservative possesses preservative capability, because the ionized portion is incapable of penetrating the microorganism. Thus, the preservative selected must be largely undissociated at the pH of the formulation being prepared. Acidic preservatives like benzoic, boric, and sorbic acids are more undissociated and thus more effective as the medium is made more acid. Conversely, alkaline preservatives are less effective in acid or neutral media and more effective in alkaline media. Thus, it is meaningless to suggest preservative effectiveness at specific concentrations unless the pH of the system is mentioned and the undissociated concentration of the agent is calculated or otherwise determined. Also, if formulative materials interfere with the solubility or availability of the preservative agent, its chemical concentration may be misleading, because it may not be a true measure of the effective concentration. Many incompatible combinations of preservative agents and other pharmaceutical adjuncts have been discovered in recent years, and undoubtedly many more will be uncovered in the future as new preservatives, pharmaceutical adjuncts, and therapeutic agents are combined for the first time. Many of the recognized incompatible combinations that inactivate the preservative contain macromolecules, including various cellulose derivatives, polyethylene glycols, and natural gums. These include tragacanth, which can attract and hold preservative agents, such as the parabens and phenolic compounds, rendering

them unavailable for their preservative function. It is essential for the research pharmacist to examine all formulative ingredients as one affects the other to ensure that each agent is free to do its job. In addition, the preservative must not interact with a container, such as a metal ointment tube or a plastic medication bottle, or with an enclosure, such as a rubber or plastic cap or liner. Such an interaction could result in decomposition of the preservative or the container closure or both, causing decomposition and contamination. Appropriate tests should be devised and conducted to prevent this type of preservative interaction.

Mode of Action

Preservatives interfere with microbial growth, multiplication, and metabolism through one or more of the following mechanisms:

- Modification of cell membrane permeability and leakage of cell constituents (partial lysis)
- Lysis and cytoplasmic leakage
- Irreversible coagulation of cytoplasmic constituents (e.g., protein precipitation)
- Inhibition of cellular metabolism, such as by interfering with enzyme systems or inhibition of cell wall synthesis
- Oxidation of cellular constituents
- Hydrolysis

A few of the commonly used pharmaceutical preservatives and their probable modes of action are presented in Table 4.6.

Preservative Utilization

Suitable substances may be added to a pharmaceutical preparation to enhance its permanency or usefulness. Such additives are suitable only if they are nontoxic and harmless in the amounts administered and do not interfere with the therapeutic efficacy or tests or assays of the preparation. Certain intravenous preparations given in large volumes as blood replenishers or as nutrients are not permitted to contain bacteriostatic additives, because the amounts required to preserve such large volumes would constitute a health hazard when administered to the patient. Thus preparations like dextrose injection, USP, and others commonly given as fluid and nutrient replenishers by intravenous injections in amounts of 500 to 1,000 mL may not contain antibacterial preservatives. On the other hand, injectable preparations

TABLE 4.6 PROBABLE MODES OF ACTION OF SOME PRESERVATIVES

PRESERVATIVE	PROBABLE MODES OF ACTION
Benzoic acid, boric acid, p-hydroxybenzoates	Denaturation of proteins
Phenols and chlorinated phenolic compounds	Lytic and denaturation action on cytoplasmic membranes and for chlorinated preservatives, also by oxidation of enzymes
Alcohols	Lytic and denaturation action on membranes
Quaternary compounds	Lytic action on membranes
Mercurials	Denaturation of enzymes by combining with thiol (-SH) groups

given in small volumes—for example, morphine sulfate injection, USP, which provides a therapeutic amount of morphine sulfate in approximately a 1-mL volume—can be preserved with a suitable preservative without the danger of the patient receiving an excessive amount of the preservative.

Examples of the preservatives and their concentrations commonly employed in pharmaceutical preparations are benzoic acid (0.1% to 0.2%), sodium benzoate (0.1% to 0.2%), alcohol (15% to 20%), phenylmercuric nitrate and acetate (0.002% to 0.01%), phenol (0.1% to 0.5%), cresol (0.1% to 0.5%), chlorobutanol (0.5%), benzalkonium chloride (0.002% to 0.01%), and combinations of methylparaben and propylparaben (0.1% to 0.2%), the latter being especially good against fungus. The required proportion

varies with the pH, dissociation, and other factors already indicated as well with the presence of other formulative ingredients with inherent preservative capabilities.

For each type of preparation to be preserved, the research pharmacist must consider the influence of the preservative on the comfort of the patient. For instance, a preservative in an ophthalmic preparation must have an extremely low degree of irritant qualities, which is characteristic of chlorobutanol, benzalkonium chloride, and phenylmercuric nitrate, frequently used in ophthalmic preparations. In all instances, the preserved preparation must be biologically tested to determine its safety and efficacy and shelf-tested to determine its stability for the intended shelf life of the product.

APPLYING THE PRINCIPLES AND CONCEPTS

GROUP ACTIVITIES

1. Develop a listing of examples where patients misunderstand the intent of the administration of a pharmaceutical dosage form.
2. Develop a listing of examples where patients misuse/abuse a pharmaceutical dosage form.
3. Explain the appropriate use of specific dosage forms for different patient types, e.g., geriatric, pediatric, visually impaired, hearing impaired.
4. Identify four ophthalmic products with differing preservative agents and provide a rationale for the selection of the specific preservative in the product.
5. Identify elixir dosage form products that contain minimal or no alcohol content. Explain the reasons for this misnomer.

INDIVIDUAL ACTIVITIES

1. Given a specific dosage form, list the signs of degradation a pharmacist might observe indicating product instability.
2. Given a concentration of drug in a liquid dosage form, determine its type of degradation rate and calculate its half life and when its concentration will be 90% of the labeled amount.
3. Compare and contrast a zero-order rate of degradation and a first-order rate of degradation.
4. Make a listing of drugs that follow a zero-order rate of degradation in a liquid dosage form.
5. Make a listing of drugs that follow first-order rates of degradation in a liquid dosage form

REFERENCES

1. Poole JW. Preformulation. FMC Corporation, 1982.
2. Brange J, Langkjaer L, Havelund S, et al. Chemical stability of insulin: Hydrolytic degradation during storage of pharmaceutical preparations. *Pharm Res* 1991;9:715–726.
3. Guideline for submitting documentation for the stability of human drugs and biologics. Rockville, MD: Food & Drug Administration, 1987.
4. FDA/ICH Regulatory Guidance on Stability. In: Federal Register, vol 63, Washington: Food & Drug Administration, 1998:9795–9843.
5. Sheinin EB. ICH Guidelines: History, Present Status, Intent. Athens, GA: International Good Manufacturing Practices Conference, 1998.
6. Rothman B. Stability is the Issue. Athens, GA: International Good Manufacturing Practices Conference, 1998.
7. General Chapter <795> Pharmaceutical Compounding-Nonsterile Preparations, and General Chapter <797> Pharmaceutical Compounding-Sterile Preparations. U.S. Pharmacopoeia 31-National Formulary 26, Rockville MD, U.S. Pharmacopoeial Convention, Inc., 2008, pp 315–336.
8. Trissel L. Trissel's Stability of Compounded Formulations. 3rd Ed. Washington: American Pharmaceutical Association, 2005.
9. Handbook of Pharmaceutical Excipients. 6th Ed. Washington: American Pharmaceutical Association, 2009.
10. Lewis R. When smell and taste go awry. *FDA Consum* 1991;25:29–33.
11. Hornstein I, Teranishi R. The chemistry of flavor. *Chem Eng News* 1967;45:92–108.
12. Murphy DH. A practical compendium on sweetening agents. *Am Pharm* 1983;NS23:32–37.
13. Jackowitz AI. Artificial sweeteners: How safe are they? *U.S. Pharmacist* 1988;13:28–31.
14. Krueger RJ, Topolewski M, Havican S. In search of the ideal sweetener. *Pharmacy Times* 1991;July:72–77.
15. Lecos CW. Sweetness minus calories = controversy. *FDA Consum* 1985;19:18–23.
16. Code of federal regulations, Title 21, Parts 70–82.
17. Colorants for drug tablets and capsules. *Drug Cosmet Ind* 1983;133(2):44.